

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07C 279/12, 257/18, C07F 5/04, C07D 211/60</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/26920 (43) International Publication Date: 3 June 1999 (03.06.99)</p>
<p>(21) International Application Number: PCT/US98/25167 (22) International Filing Date: 23 November 1998 (23.11.98) (30) Priority Data: 08/980,114 26 November 1997 (26.11.97) US 08/979,440 26 November 1997 (26.11.97) US (71) Applicant (for all designated States except US): CORVAS INTERNATIONAL, INC. [US/US]; 3030 Science Park Road, San Diego, CA 92121 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SEMPLE, Joseph, Edward [US/US]; 9711 Caminito Pudregal, San Diego, CA 92131 (US). LIM-WILBY, Marguerita, S. [US/US]; 5749 Caminito Empresa, La Jolla, CA 92037 (US). BRUNCK, Terence, K. [US/US]; 4949 Quincy Street, San Diego, CA 92109 (US). (74) Agents: BIGGS, Suzanne, L. et al.; Lyon &amp; Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: SUBSTITUTED 3-AMINO-2-HYDROXYPHENYLACETAMIDE DERIVATIVES AS ENZYME INHIBITORS (II) (57) Abstract  The present invention provides peptide aldehydes having an 3-amino-2-hydroxyphenyl acetamide group as part of the peptide backbone and an arginine group or analog at P1. These compounds are potent and specific or inhibitors of thrombin. Their pharmaceutically acceptable salts, pharmaceutically acceptable compositions thereof, and methods of using them as therapeutic agents for disease states in mammals characterized by abnormal thrombosis are also described. Also described are 3-amino-2-hydroxyphenyl-acetamide derivatives having in history activity towards proteases of the trypsin/chymotrypsin class.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	ME	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5                   SUBSTITUTED 3-AMINO-2-HYDROXYPHENYLACETAMIDE DERIVATIVES AS  
                  ENZYME INHIBITORS (II)

Technical Field

          In one aspect, the present invention relates to compounds which are potent and specific inhibitors of thrombin. In  
10 another aspect, the present invention relates to novel peptide aldehydes, their pharmaceutically acceptable salts, and  
pharmaceutically acceptable compositions thereof which are  
useful as potent and specific inhibitors of blood coagulation  
15 in vitro and in vivo in mammals. In yet another aspect, the  
invention relates to methods of using these inhibitors as  
therapeutic agents for disease states in mammals characterized  
by abnormal thrombosis.

20 Background

          Normal hemostasis is the result of a complex balance between the processes of clot formation (blood coagulation) and clot dissolution (fibrinolysis). The complex interactions between blood cells, specific plasma proteins and the vascular  
25 surface, maintain the fluidity of blood unless injury and blood loss occur.

          Blood coagulation is the culmination of a series of amplified reactions in which several specific zymogens of serine proteases in plasma are activated by limited  
30 proteolysis. Nemerson, Y. and Nossel, H.L., Ann. Rev. Med., 33: 479 (1982). This series of reactions results in the formation of an insoluble fibrin matrix which is required for the stabilization of the primary hemostatic plug. The interaction and propagation of the activation reactions occurs

through the extrinsic and intrinsic pathways of coagulation.

These pathways are highly inter-dependent and converge in the formation of the serine protease, Factor Xa. Factor Xa catalyzes the penultimate step in the blood coagulation cascade which is the formation of the serine protease thrombin. This step occurs following the assembly of the prothrombinase complex which is composed of factor Xa, the non-enzymatic co-factor Va and the substrate prothrombin assembled on the surface of adhered, activated platelets or systemically circulating membranous microparticles.

Proteolytic activation of zymogen factor X to its catalytically active form, factor Xa, can occur by either the intrinsic or extrinsic coagulation pathways.

The intrinsic pathway is referred to as "intrinsic" because everything needed for clotting is in the blood. Saito, H., "Normal Hemostatic Mechanisms", Disorders of Hemostasis, pp. 27-29, Grune & Stratton, Inc. (O. D. Ratnoff, M.D. and C. D. Forbes, M.D. edit. 1984). This pathway is comprised of the zymogen serine proteases, factors IX and XI, and the non-enzymatic co-factor, factor VIII. The initiation of the intrinsic pathway results in the activation of factor XI to XIa. Factor XIa catalyzes the activation of factor IX to factor IXa which in combination with the activated form of factor VIII on an appropriate phospholipid surface, results in the formation of the tenase complex. This complex also catalyzes the formation of the serine protease, factor Xa, from its zymogen, factor X which subsequently results in clot formation.

The extrinsic pathway is referred to as "extrinsic" because the tissue factor which binds to and facilitates the activation of factor VII comes from outside the blood. Saito, Id. The major components of this pathway are the zymogen serine protease, factor VII, and the membrane bound protein, tissue factor. The latter serves as the requisite non-enzymatic co-factor for this enzyme. The initiation of this pathway is thought to be an autocatalytic event resulting from

the activation of zymogen factor VII by trace levels of activated factor VII (factor VIIa), both of which are bound to newly exposed tissue factor on membrane surfaces at sites of vascular damage. The factor VIIa/tissue factor complex  
5 directly catalyzes the formation of the serine protease, factor Xa, from its zymogen, factor X. Exposure of blood to injured tissue initiates blood clotting by the extrinsic pathway.

The formation of thrombin is catalyzed by factor Xa  
10 following the assembly of the catalytic prothrombinase complex as reviewed by Mann, K. G. et. al., "Surface-Dependent Reactions of the Vitamin K-Dependent Enzyme Complexes", Blood, 76: 1-16 (1990). This complex is composed of factor Xa, the non-enzymatic co-factor Va and the substrate prothrombin all  
15 assembled on an appropriate phospholipid surface. The requirement of a macromolecular complex for efficient catalysis results in the protection of factor Xa from natural anticoagulant mechanisms such as heparin-antithrombin III mediated inhibition. Teite, J. M. and Rosenberg, R. D.,  
20 "Protection of Factor Xa from neutralization by the heparin-antithrombin complex", J. Clin. Invest., 71: 1383-1391(1983). In addition, sequestration of factor Xa in the prothrombinase complex also renders it resistant to inhibition by exogenous heparin therapy which also requires antithrombin III to elicit  
25 its anticoagulant effect.

Thrombin is the primary mediator of thrombus formation. Thrombin acts directly to cause formation of insoluble fibrin from circulating fibrinogen. In addition, thrombin activates the zymogen factor XIII to the active transglutaminase factor  
30 XIIIa which acts to covalently stabilize the growing thrombus by crosslinking the fibrin strands. Lorand, L. and Konishi, K., Arch. Biochem. Biophys., 105: 58 (1964). Beyond its direct role in the formation and stabilization of fibrin rich clots, the enzyme has been reported to have profound  
35 bioregulatory effects on a number of cellular components within the vasculature and blood. Shuman, M.A., Ann. NY Acad.

Sci., 405: 349 (1986).

It is believed that thrombin is the most potent agonist of platelet activation, and it has been demonstrated to be the primary pathophysiologic-mediator of platelet-dependent arterial thrombus formation. Edit, J.F. et al., J. Clin. Invest., 84: 18 (1989). Thrombin-mediated platelet activation leads to ligand-induced inter-platelet aggregation principally due to the bivalent interactions between adhesive ligands such as fibrinogen and fibronectin with platelet integrin receptors such as glycoprotein IIb/IIIa which assume their active conformation following thrombin activation. Berndt, M.C. and Phillips, D.R., Platelets in Biology and Pathology, pp 43-74, Elsevier/North Holland Biomedical Press (Gordon, J.L. edit. 1981). Thrombin-activated platelets can also support further thrombin production through the assembly of new prothrombinase and tenase (factor IXa, factor VIIIa and factor X) catalytic complexes on the membrane surface of intact activated platelets and platelet-derived microparticles, following thrombin-mediated activation of the non-enzymatic cofactors V and VIII, respectively. Tans, G. et al., Blood, 77: 2641 (1991). This positive feedback process results in the local generation of large concentrations of thrombin within the vicinity of the thrombus which supports further thrombus growth and extension. Mann, K.G. et al., Blood, 76: 1 (1990).

In contrast to its prothrombotic effects, thrombin has been shown to influence other aspects of hemostasis. These include its effect as an important physiological anticoagulant. The anticoagulant effect of thrombin is expressed following binding of thrombin to the endothelial cell membrane glycoprotein, thrombomodulin. This is thought to result in an alteration of the substrate specificity of thrombin thereby allowing it to recognize and proteolytically activate circulating protein C to give activated protein C (aPC). Musci, G. et al., Biochemistry, 27: 769 (1988). aPC is a serine protease which selectively inactivates the non-enzymatic co-factors Va and VIIIa resulting in a down-

regulation of thrombin formation by the prothrombinase and tenase catalytic complexes, respectively. Esmon, C.T., *Science*, **235**: 1348 (1987). The activation of protein C by thrombin in the absence of thrombomodulin is poor.

5. Thrombin has also been shown to be a potent direct mitogen for a number of cell types, including cells of mesenchymal origin such as vascular smooth muscle cells. Chen, L.B. and Buchanan, J.M., *Proc. Natl. Acad. Sci. USA*, **72**: 131 (1975). The direct interaction of thrombin with vascular smooth muscle also results in vasoconstriction. Walz, D.A. et al., *Proc. Soc. Exptl. Biol. Med.*, **180**: 518 (1985). Thrombin acts as a direct secretagogue inducing the release of a number of bioactive substances from vascular endothelial cells including tissue plasminogen activator. Levin, E.G. et al., *Thromb. Haemost.*, **56**: 115 (1986). In addition to these direct effects on vascular cells, the enzyme can indirectly elaborate potent mitogenic activity on vascular smooth muscle cells by the release of several potent growth factors (e.g. platelet-derived growth factor and epidermal growth factor) from platelet  $\alpha$ -granules following thrombin-induced activation. Ross, R., *N. Engl. J. Med.*, **314**: 408 (1986).

Many significant disease states are related to abnormal hemostasis. With respect to the coronary arterial vasculature, abnormal thrombus formation due to the rupture of an established atherosclerotic plaque is the major cause of acute myocardial infarction and unstable angina. Moreover, treatment of an occlusive coronary thrombus by either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA) is often accompanied by an acute thrombotic reclosure of the affected vessel which requires immediate resolution. With respect to the venous vasculature, a high percentage of patients undergoing major surgery in the lower extremities or the abdominal area suffer from thrombus formation in the venous vasculature which can result in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Disseminated

intravascular coagulopathy commonly occurs within both vascular systems during septic shock, certain viral infections and cancer and is characterized by the rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the vasculature leading to widespread organ failure.

Pathogenic thrombosis in the arterial vasculature is a major clinical concern in today's medicine. It is the leading cause of acute myocardial infarction which is one of the leading causes of death in the western world. Recurrent arterial thrombosis also remains one of the leading causes of failure following enzymatic or mechanical recanalization of occluded coronary vessels using thrombolytic agents or percutaneous transluminal coronary angioplasty (PTCA), respectively. Ross, A.M., Thrombosis in Cardiovascular Disorder, p. 327, W.B. Saunders Co. (Fuster, V. and Verstraete, M. edit. 1991); Califf, R.M. and Willerson, J.T., Id. at p 389. In contrast to thrombotic events in the venous vasculature, arterial thrombosis is the result of a complex interaction between fibrin formation resulting from the blood coagulation cascade and cellular components, particularly platelets, which make up a large percentage of arterial thrombi. Heparin, the most widely used clinical anticoagulant administered i.v., has not been shown to be universally effective in the treatment or prevention of acute arterial thrombosis or rethrombosis. Prins, M.H. and Hirsh, J., J. Am. Coll. Cardiol., 67: 3A (1991).

Besides the unpredictable, recurrent thrombotic reocclusion which commonly occurs following PTCA, a profound restenosis of the recanalized vessel occurs in 30 to 40% of patients 1 to 6 months following this procedure. Califf, R.M. et al., J. Am. Coll. Cardiol., 17: 2B (1991). These patients require further treatment with either a repeat PTCA or coronary artery bypass surgery to relieve the newly formed stenosis. Restenosis of a mechanically damaged vessel is not a thrombotic process but instead is the result of a



hyperproliferative response in the surrounding smooth muscle cells which over time results in a decreased luminal diameter of the affected vessel due to increased muscle mass. Id. As for arterial thrombosis, there is currently no effective  
5 pharmacologic treatment for the prevention of vascular restenosis following mechanical recanalization.

The need for safe and effective therapeutic anticoagulants has in one aspect focused on the role of the serine protease thrombin in blood coagulation.

10 Most preferred natural substrates for thrombin are reported to contain an uncharged amino acid in the P3 recognition subsite. For example, the thrombin cleavage site on the Aa chain of fibrinogen, which is the primary  
15 physiological substrate for thrombin, is reported to contain a glycine residue in this position while the cleavage site on the Bb chain contains a serine, as shown below:

P4	P3	P2	P1	P1
Gly-Gly-Val-Arg/Gly Fibrinogen Aa Chain				
[SEQ. ID. NO. 1]				
Phe-Ser-Ala-Arg/Gly Fibrinogen Bb Chain				
[SEQ. ID. NO. 2]				

Peptidyl derivatives having an uncharged residue in the P3 position are said to bind to the active site of thrombin  
25 and thereby inhibit the conversion of fibrinogen to fibrin and cellular activation have been reported. These derivatives have either an aldehyde, chloromethyl ketone or boronic acid functionality associated with the P1 amino acid. For example, substrate-like peptidyl derivatives such as D-phenylalanyl-prolyl-argininal (D-Phe-Pro-Arg-al), D-phenylalanyl-prolyl-arginine-chloromethyl ketone (P-PACK) and acetyl-D-phenylalanyl-prolyl-boroarginine (Ac-(D-Phe)-Pro-boroArg) have  
30 been reported to inhibit thrombin by directly binding to the active site of the enzyme. Bajusz, S., Symposia Biologica Hungarica, 25: 277 (1984), Bajusz, S. et al., J. Med. Chem., 33: 1729 (1990) and Bajusz, S. et al., Int. J. Peptide Protein  
35

Res. 12: 217 (1970); Kettner, C. and Shaw, E., Methods Enzymol., 80: 826 (1987), Kettner, C. et al., EP 293,881 (published December 7, 1988), Kettner, C., et al., J. Biol. Chem., 265: 18209 (1990). These molecules have been reported  
5 to be potent anticoagulants in the prevention of platelet-rich arterial thrombosis. Kelly, A.B. et al., Thromb. Haemostas., 65: 736 at abstract 257 (1991). Other peptidyl aldehydes have been proposed or reported as inhibitors of thrombin. Bey, P. et al., EP 363,284 (published April 11, 1990) and  
10 Balasubramanian, N. et al., EP 526,877 (published February 10, 1993).

Peptidyl compounds which are said to be active site inhibitors of thrombin but which differ in structure from those containing a uncharged amino acid in the P3 recognition  
15 subsite have been reported.

The compound, Argatroban (also called 2R,4R-4-methyl-1-[N-2-(3-methyl-1,2,3,4-tetrahydro-8-quinolinesulfonyl)-L-argininy]-2-piperidinecarboxylic acid), is also reported to bind directly to the active site of thrombin and has been  
20 thought to be the most potent and selective compound in the class of non-peptidyl inhibitors of this enzyme. Okamoto, S. et al., Biochem. Biophys. Res. Commun., 101: 440 (1981). Argatroban has been reported to be a potent antithrombotic agent in several experimental models of acute arterial  
25 thrombosis. Jang, I.K. et al., in both Circulation, 81: 219 (1990) and Circ. Res., 67: 1552 (1990).

Peptidyl compounds which are said to be inhibitors of thrombin and whose mode of action is thought to be by binding to both the active site and another site on the enzyme have  
30 been reported. Hirudin and certain peptidyl derivatives of hirudin have been reported to inhibit both conversion of fibrinogen to fibrin and platelet activation by binding to either both the active site and exo site, or the exo site only, of thrombin. Markwardt, F., Thromb. Haemostas., 66: 141  
35 (1991). Hirudin is reported to be a 65 amino acid polypeptide originally isolated from leech salivary gland extracts. It is

- said to be one of the most potent inhibitors of thrombin known. Marki, W.E. and Wallis, R.B., Thromb. Haemostas., **64**: 344 (1990). It has been reported to inhibit thrombin by binding to both its anion-binding exo-site and to its catalytic active site which are distinct and physically distant from each other. Rydel, T.J. et al., Science, **249**:277 (1990). Hirudin has been reported to be a potent antithrombotic agent in vivo. Markwardt, F. et al., Pharmazie, **43**: 202 (1988); Kelly, A.B. et al., Blood, **77**: (1991). In addition to its antithrombotic effects, hirudin has been reported to also effectively inhibit smooth muscle proliferation and the associated restenosis following mechanical damage to a atherosclerotic rabbit femoral artery. Sarembock, I.J. et al., Circulation, **84**: 232 (1991).
- 15 Hirugen has been reported to be a peptide derived from the anionic carboxy-terminus of hirudin. It is reported to bind only to the anion binding exo-site of thrombin and thereby inhibit the formation of fibrin but not the catalytic turnover of small synthetic substrates which have access to the unblocked active site of the enzyme. Maraganore, J.M. et al., J. Biol. Chem., **264**: 8692 (1989); Naski, M.C. et al., J. Biol. Chem., **265**: 13484 (1990). The region of hirudin represented by hirugen has been reported, as according to by x-ray crystallographic analysis, to bind directly to the exo site of thrombin. Skrzypczak-Jankun, E. et al., Thromb. Haemostas., **65**: 830 at abstract 507 (1991). Moreover, the binding of hirugen has also been reported to enhance the catalytic turnover of certain small synthetic substrates by thrombin, indicating that a conformational change in the enzyme active site may accompany occupancy of the exo-site. Liu, L.W. et al., J. Biol. Chem., **266**:16977 (1991). Hirugen also is reported to block thrombin-mediated platelet aggregation. Jakubowski, J.A. and Maraganore, J.M., Blood, **75**: 399 (1990).
- 35 A group of synthetic chimeric molecules comprised of a hirugen-like sequence linked by a glycine-spacer region to the

peptide, D-phenylalanyl-pr lyl-arginine, which is based on a preferred substrate recognition site for thrombin, has been termed to be hirulog. Maraganore et al., U.S. Patent No. 5,196,404 (March 23, 1993). The hirugen-like sequence is said to be linked to this peptide through the C-terminal end of the peptide. Maraganone, J.M. et al., Biochemistry, 29: 7095 (1990). The hirulogs have been reported to be an effective antithrombotic agents in preventing both fibrin-rich and platelet-rich thrombosis. Maraganone, J.M. et al., Thromb. Haemostas., 65: 651 at abstract 17 (1991).

Certain benzamidines have been reported to inhibit thrombin though non-selectively. 4-amidinophenylpyruvic acid (APPA) has been reported to be a thrombin inhibitor with low toxicity and favourable pharmacokinetics. However, this compound was reported to be non-selective, inhibiting trypsin, plasmin and kallikrein. Markwardt et al., Thromb. Res., 1:243-52 (1972). Other benzamidine-derived structures which have been reported to inhibit thrombin include the cyclic amides of N<sup>a</sup>-substituted 4-amidinophenylalanine and 2-amino-5-(4-amidinophenyl)-1-valeric acid. The inhibitory constant displayed by these compounds was reported to be in the micromolar range. Markwardt et al., Thromb. Res., 17:425-31 (1980). Moreover, derivatives of 4-amidinophenylalanine whose  $\alpha$ -amino group is linked to the arylsulfonyl residue via an  $\omega$ -aminoalkylcarboxylic acid as spacer have also been assessed for their inhibitory effect. Among these N<sup>a</sup>-(2-naphthylsulphonylglycyl)-4-amidinophenylalanine piperidide ( $\alpha$ -NAPAP) has been reported to possess an affinity for thrombin ( $K_i=6 \times 10^{-9}$  M). Banner et al., J. Biol. Chem., 266:20085 (1991) and Sturzebecher et al., Thromb. Res., 29:635-42 (1983).

Certain bis-benzamidines have been reported to inhibit thrombin. The antithrombin activity of bis-benzamidines was reported to increase with the length and bulkiness of the central chain. However, these compounds were reported to be generally toxic in the micromolar range where they are also

inhibitory. Geratz et. al., Thromb. Diath. Haemorrh.,  
29:154-67 (1973); Geratz et. al., J. Med. Chem., 16:970-5  
(1973); Geratz et. al., J. Med. Chem., 19:634-9 (1976);  
Walsmann et. al., Acta Biol. Med. Germ., 35:K1-8 (1976); and  
5 Hauptmann et. al., Acta Biol. Med. Germ., 35:635-44 (1976).


Certain amidino-bearing aromatic ring structures such as  
b-naphthamidines have been reported to possess modest  
antithrombin and anticoagulant activity. This class of  
compounds include the non-selective 6-amidino-2-naphthyl- 4-  
10 guanidinobenzoate dimethanesulfonate (FUT 175). Fuji et al.,  
Biochim. Biophys. Acta, 661:342-5 (1981); and Hitomi et. al.,  
Haemostasis, 15:164-8 (1985).

Certain phenylguanidines have been reported to inhibit  
thrombin. Derivatives of 4-guanidinophenylalanine with  
15 inhibitory constants in the micromolar range have been  
reported to inhibit thrombin. This class includes the Na<sup>+</sup>-  
tosylated and dansylated 4-guanidino phenylalanine  
piperidides. Claeson et. al., Thromb. Haemostas., 50:53  
(1983). Another compound, [ethyl p-(6-guanidinohexanoyloxy)  
20 benzoate] methane sulfonate (FOY) was reported to be a non-  
selective competitive inhibitor of thrombin. Ohno et al.,  
Thromb. Res., 19:579-588 (1980).

Certain compounds having inhibitory activity toward  
serine proteases, including thrombin, factor Xa, and trypsin,  
25 are disclosed within the following commonly assigned United  
States patents or published PCT applications: US 5,371,072;  
5,492,895; 5,534,498; 5,597,804; 5,637,599; 5,646,165;  
5,656,600; 5,656,645; WO 94/13693; WO 95/35311; WO 95/35313;  
WO 96/19493.

Summary of the Invention

Novel compounds of the present invention include compounds of the formula (I):



wherein

(a) X is selected from the group consisting of  $-S(O)_2-$ ,  $-N(R')-S(O)_2-$ ,  $-C(=O)-$ ,  $-OC(=O)-$ ,  $-NH-C(=O)-$ ,  $-P(O)(R'')$  and a direct link, wherein R' is hydrogen, alkyl of 1 to about 4 carbon atoms, aryl of about 6 to about 14 carbon atoms or aralkyl of about 6 to about 16 carbon atoms, and R'' is  $NHR'$ ,  $OR'$ ,  $R'$ , or  $SR'$ ;

(b) R<sub>1</sub> is selected from the group consisting of:

(1) alkyl of 1 to about 12 carbon atoms,  
 (2) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 5 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,

(3) cycloalkyl of 3 to about 15 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,

(4) heterocycloalkyl of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and  $S(O)_i$ , wherein i is 0, 1 or 2, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub>

to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

(5) heterocyclo of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>; and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl or alkoxy, or -CO<sub>2</sub>R',

(6) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

(7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(8) heteroaryl of about 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(9) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,



(10) heteroaralkyl of 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(11) aralkenyl of about 8 to about 16 carbon atoms having 5 to about 14 ring atoms which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,



(12) heteroaralkenyl of 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms,

wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

5



10



15

(17) perfluoroalkyl of 1 to about 12 carbon atoms,

(18) perfluoroaryl of about 6 to about 14 carbon

atoms,

(19) perfluoroaralkyl of about 7 to about 15 carbon

atoms,

(20) hydrogen, and

20

(21) , wherein is a 5 to 7 membered heterocyclic ring having 3 to 6 ring carbon atoms, where V is -CH<sub>2</sub>-, -O-, -S(=O)-, -S(O)<sub>2</sub>- or -S-,

wherein Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>3</sub> are

(i) independently selected from the group  
25 consisting of hydrogen, halogen, cyano, nitro, tetrazolyl,




- amin, guanidino, amidino, methylamino, methylguanidin,  
 $-\text{CF}_3$ ,  $-\text{CF}_2\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{CH}(\text{CF}_3)_2$ ,  $-\text{C}(\text{OH})(\text{CF}_3)_2$ ,  $-\text{OCF}_3$ ,  
 $-\text{OCF}_2\text{CF}_3$ ,  $-\text{OC}(\text{O})\text{NH}_2$ ,  $-\text{OC}(\text{O})\text{NHZ}_1$ ,  $-\text{OC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{NHC}(\text{O})\text{Z}_1$ ,  
 $-\text{NHC}(\text{O})\text{NH}_2$ ,  $-\text{NHC}(\text{O})\text{NHZ}_1$ ,  $-\text{NHC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{NH}_2$ ,  
 5.  $-\text{C}(\text{O})\text{NHZ}_1$ ,  $-\text{C}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OZ}_1$ ,  $-\text{P}(\text{O})_3\text{H}$ ,  $-\text{P}(\text{O})_3\text{H}_2$ ,  $-\text{PH}(\text{O})\text{OH}$ ,  
 $-\text{P}(\text{O})_3(\text{Z}_1)_2$ ,  $-\text{S}(\text{O})_3\text{H}$ ,  $-\text{S}(\text{O})_m\text{Z}_1$ ,  $-\text{Z}_1$ ,  $-\text{OZ}_1$ ,  $-\text{OH}$ ,  $-\text{NH}_2$ ,  $-\text{NHZ}_1$ ,  
 and  $-\text{NZ}_1\text{Z}_2$ , wherein  $m$  is 0, 1 or 2, and  $\text{Z}_1$  and  $\text{Z}_2$  are

independently selected from the group consisting of alkyl of  
 1 to about 12 carbon atoms, aryl of about 6 to about 14  
 10 carbon atoms, heteroaryl of about 5 to about 14 atoms having  
 1 to about 9 carbon atoms, aralkyl of about 7 to about 15  
 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms  
 having about 3 to about 9 carbon atoms, or


- (ii)  $\text{Y}_1$  and  $\text{Y}_2$  are selected together to be  
 15  $-\text{O}[\text{C}(\text{Z}_3)(\text{Z}_4)]_q\text{O}-$ , wherein  $q$  is an integer from 1 to 4 and  $\text{Z}_3$   
 and  $\text{Z}_4$  are independently selected from the group consisting  
 of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of  
 about 6 to about 14 carbon atoms heteroaryl of about 5 to  
 about 14 atoms having 1 to about 9 carbon atoms, aralkyl of  
 20 about 7 to about 15 carbon atoms, and heteroaralkyl of about  
 6 to about 11 atoms having about 3 to about 9 carbon atoms;



(c)  $\text{R}_2$  is selected from the group consisting of  
 hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of  
 about 2 to about 4 carbon atoms;

- 25 (d)  $\text{R}_3$  is selected from the group consisting of



where  $d$  is an integer from 0 to 5 and  $W$  is  $-\text{N}-$  or  $-\text{CH}-$ ;

- (e)  $\text{R}_4$  is selected from the group consisting of  
 (i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein  $n$  is 0, 1  
 30 or 2, and  $\text{R}_1$  is as defined above, with the proviso that  $\text{R}_1$  is  
 not a camphor derivative, or  heterocycle,

- (ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ ,  
(iii) halogen,  
(iv) alkyl of 1 to 6 carbon atoms substituted with  
Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of  
5 hydrogen,  $-\text{OH}$ ,  $-\text{OR}_8$ ,  $-\text{CH}_3$ , halogen,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{OR}_8$  and  
 $-\text{S}(\text{O})_p\text{R}_8$ , wherein R<sub>8</sub> is alkyl of 1 to about 6 carbon atoms,  
and p is 0, 1 or 2  
(v) alkyl of 1 to 3 carbon atoms substituted with  
cycloalkyl of 3 to 5 carbon atoms,  
10 (vi) alkenyl of about 3 to about 6 carbon atoms,  
(vii) cycloalkyl of about 3 to about 10 carbon  
atoms,  
(viii) heteroaryl of about 5 to about 10 ring atoms  
having 1 to about 9 ring carbon atoms and the remainder of  
15 the ring atoms heteroatoms selected from oxygen, sulfur and  
nitrogen, and  
(ix) heteroaralkyl of about 5 to about 10 ring  
atoms having about 2 to about 15 carbon atoms which include 1  
to about 9 ring carbon atoms and the remainder of the ring  
20 atoms heteroatoms selected from oxygen, sulfur and nitrogen;
- (f) R<sub>5</sub> is selected from the group consisting of  
(i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein R<sub>1</sub> is as  
defined above, with the proviso that R<sub>1</sub> is not a camphor  
25 derivative or  heterocycle,  
(ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ , and  
(iii) halogen;  
(g) R<sub>6</sub> is selected from the group consisting of  
(i)  $\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein R<sub>1</sub> is as  
30 defined above, with the proviso that R<sub>1</sub> is not a camphor  
derivative or  heterocycle,  
(ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ ,  
(iii) halogen,  
(iv) alkyl of 1 to about 12 carbon atoms

substituted with Z<sub>6</sub>, wherein Z<sub>6</sub> is selected from the group consisting of hydrogen, hydroxy, halogen, -OR<sub>9</sub>, -NHR<sub>9</sub>, -C(O)OH, -C(O)OR<sub>9</sub>, and -S(O)<sub>p</sub>R<sub>9</sub>, wherein R<sub>9</sub> is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon atoms, aralkyl of about 7 to about 12 carbon atoms, heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted in the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>; or heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, and which is optionally mono-, di- or tri- substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>;

(v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms;

(vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 5 to about 10 carbon atoms or heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms,

(vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(ix) aralkenyl of 6 to about 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

(x) heteroaralkenyl of about 5 to about 10 ring

atoms having 1 to 9 ring carbon atoms and having the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, as defined above; and

(h) R<sub>7</sub> is selected from the group consisting of

(i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above, with the

proviso that R<sub>1</sub> is not a camphor derivative or  heterocycle,

(ii) -CF<sub>3</sub>, or -CF<sub>2</sub>H,

(iii) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(iv) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>,


(vi) heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>, and

(vii) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>, and wherein each of R<sub>10</sub> and R<sub>11</sub> is independently selected and is alkyl of 1 to about 4 carbon atoms or hydrogen; or pharmaceutically acceptable salts thereof.

Among other factors, the present invention is based on our finding that these novel compounds are active as selective inhibitors of thrombin *in vivo* and *in vitro*. Furthermore, certain of the preferred compounds of the present invention have been found to exhibit advantageous

selectivity in that they are potent inhibitors of thrombin, but are much less active and much less potent in inhibiting plasmin and trypsin. Certain compounds of the present invention may also have broader serine protease inhibitory activity.

According to an alternate aspect of the present invention, also provided are compounds of formula (II) which have inhibitory activity towards proteases of the trypsin/chymotrypsin class. This aspect of the invention is directed to compounds of formula (II):



(II)

wherein a substituted phenol or phenyl ether is present in the compound in place of the amide bond of an inhibitor that makes hydrogen bonds to amino acid 216 in the trypsin/chymotrypsin class of serine proteases. Thus, according to this aspect of the present invention, provided are compounds of formula (II), wherein X, R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are as defined hereinabove in connection with formula (I), and

- (a) R<sub>12</sub> is selected from the group consisting of
- (i) alkyl of 1 to about 12 carbon atoms optionally substituted with amino, amidino, or guanidino,
  - (ii) aralkyl of 7 to about 15 carbon atoms which is optionally substituted on a ring carbon with amino, amidino or guanidino,
  - (iii) aryl of 6 to about 14 carbon atoms which is optionally substituted on a ring carbon with amino, amidino, or guanidino, and
  - (iv) heteroaralkyl of 5 to about 14 ring atoms having about 1 to about 13 ring carbon atoms and the

remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur which is optionally substituted on a ring carbon with amino, amidino or guanidino;

and

5 (b) TS is selected from the group consisting of

- (i) -CHO,
- (ii) -C(O)CF<sub>3</sub>,
- (iii) -C(O)CF<sub>2</sub>CF<sub>3</sub>,
- (iv) -C(O)-C(O)-NHR<sub>13</sub>,
- 10 (v) -C(O)-C(O)<sub>2</sub>R<sub>13</sub>,
- (vi) -C(O)-C(O)-NR<sub>13</sub>R<sub>14</sub>,
- (vii) -B(OR<sub>13</sub>)(OR<sub>14</sub>),
- (viii) -C(O)CH<sub>2</sub>Cl, and
- 15 (ix) -C(O)R<sub>15</sub>;

15 wherein R<sub>13</sub> and R<sub>14</sub> are independently selected from hydrogen lower alkyl of 1 to about 6 carbon atoms, cycloalkyl of 3 to about 10 carbon atoms, aralkyl of 7 to about 15 carbon atoms, and heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9  
20 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur, and R<sub>15</sub> is selected from aryl of 6 to about 14 carbon atoms and heteroaryl of 5 to about 14 ring atoms having 1 to about 13  
25 ring carbon atoms and the remainder of the ring atoms selected from oxygen, nitrogen and sulfur. In the case where TS is (vii), additionally -OR<sub>13</sub> and -OR<sub>14</sub> may be taken together to be -O[C(Z<sub>3</sub>)(Z<sub>4</sub>)]<sub>q</sub>O- wherein Z<sub>3</sub>, Z<sub>4</sub> and q are as defined in connection with formula (I). Also included are  
30 pharmaceutically acceptable salts of the compounds of formula (II).

In another aspect, the present invention is directed to pharmaceutical compositions comprising a therapeutically effective amount of a compound of the present invention and a pharmaceutically acceptable carrier.

In yet another aspect, the present invention is directed to methods of using the compounds and pharmaceutical compositions of formula (I) for the prevention of or treating of thrombosis in a mammal suspected of having a condition of abnormal thrombosis, which comprises administering to said  
5 mammal a therapeutically effective amount of a compound of the present invention or pharmaceutical composition comprising such a compound.

#### 10 Definitions

In accordance with the present invention and as used herein, the following terms are defined to have following meanings, unless explicitly stated otherwise:

The term "alkenyl" refers to unsaturated aliphatic  
15 groups having at least one double bond.

The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain and cyclic groups.

The terms "alkoxy" and "alkoxyl" refer to a group having the formula, R-O-, wherein R is an alkyl group.

20 The term "alkoxycarbonyl" refers to -C(O)OR wherein R is alkyl.

The term "aralkenyl" refers to an alkenyl group substituted with an aryl group.

The term "aralkyl" refers to an alkyl group substituted  
25 with an aryl group. Suitable aralkyl groups include benzyl, phenethyl, and the like, all of which may be optionally substituted.

The term "aryl" refers to aromatic groups which have at least one ring having a conjugated pi electron system and  
30 includes carbocyclic aryl, heterocyclic aryl and biaryl groups, all of which may be optionally substituted.

The term "aryloxy" refers to a group having the formula, R-O-, wherein R is an aryl group.

The term "aralkoxy" refers to a group having the formula,  
35 R-O-, wherein R is an aralkyl group.

The term "amino acid" refers to natural amino acids,


unnatural amino acids, and amino acid analogs, all in their D and L stereoisomers if their structure allow such stereoisomeric. Natural amino acids include alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp),  
5 cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), tryptophan (Trp), tyrosine (Tyr) and valine (Val). Unnatural amino acids  
10 include, but are not limited to azetidinecarboxylic acid, 2-aminoadipic acid, 3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid,  
15 2-aminopimelic acid, 2,4-diaminoisobutyric acid, desmosine, 2,2'-diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylglycine, N-  
20 methylisoleucine, N-methylvaline, norvaline, norleucine, ornithine and pipercolic acid. Amino acid analogs include the natural and unnatural amino acids which are chemically blocked, reversibly or irreversibly, or modified on their N-terminal amino group or their side-chain groups, as for  
25 example, methionine sulfoxide, methionine sulfone, S-(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine sulfoxide and S-(carboxymethyl)-cysteine sulfone.


The term "amino acid analog" refers to an amino acid wherein either the C-terminal carboxy group, the N-terminal  
30 amino group or side-chain functional group has been chemically modified to another functional group. For example, aspartic acid-(beta-methyl ester) is an amino acid analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid  
35 analog of alanine.

The term "amino acid residue" refers to radicals having



the structure: (1)  $-C(O)-R-NH-$ , wherein R typically is  $-CH(R')-$ , wherein R' is H or a carbon containing substituent;



or (2) , wherein p is 1, 2 or 3 representing the azetidinedicarboxylic acid, proline or pipecolic acid residues, respectively.

"Biaryl" refers to phenyl substituted by carbocyclic or heterocyclic aryl as defined herein, ortho, meta or para to the point of attachment of the phenyl ring.

"Brine" refers to an aqueous saturated solution of sodium chloride.

"Carbocyclic aryl" refers to aromatic groups wherein the ring atoms on the aromatic ring are carbon atoms.

Carbocyclic aryl groups include monocyclic carbocyclic aryl groups and naphthyl groups, all of which may be optionally substituted. Suitable carbocyclic aryl groups include phenyl and naphthyl. Suitable substituted carbocyclic aryl groups include indene and phenyl substituted by one to two substituents such being advantageously lower alkyl, hydroxy, lower alkoxy, lower alkoxycarbonyl, halogen, trifluoromethyl, difluoromethyl, nitro, and cyano. Substituted naphthyl refers to naphthyl, more preferably 1- or 2-naphthyl, substituted by  $Y_1$ ,  $Y_2$  and/or  $Y_3$  as defined in connection with formula (I) hereinabove..

"Cycloalkenyl" refers to a cyclic alkenyl group. Suitable cycloalkenyl groups include, for example, cyclopentenyl and cyclohexenyl.

"Cycloalkyl" refers to a cyclic alkyl group. Suitable cycloalkyl groups include, for example, cyclohexyl, cyclopropyl, cyclopentyl, and cycloheptyl.

"Cyclohexylmethyl" refers to a cyclohexyl group attached to  $CH_2$ .

The term "halogen" refers to fluorine, chlorine, bromine and iodine.

"Heteroaralkenyl" refers to an alkenyl group substituted with a heteroaryl, and includes those heterocyclic systems described in "Handbook of Chemistry and Physics", 49th edition, 1968, R.C. Weast, editor; The Chemical Rubber Co., Cleveland, OH. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems. Preferably the alkenyl group has from 2 to about 6 carbon of atoms.

"Heteroaralkyl" refers to an alkyl group substituted with a heteroaryl, such as picolyl, and includes those heterocyclic systems described in "Handbook of Chemistry and Physics", 49th edition, 1968, R.C. Weast, editor; The Chemical Rubber Co., Cleveland, OH. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems. Preferably the alkyl group has from 1 to about 6 carbon atoms.

"Heteroaryl" refers to aryl groups having from 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms, and includes those heterocyclic systems described in "Handbook of Chemistry and Physics", 49th edition, 1968, R.C. Weast, editor; The Chemical Rubber Co., Cleveland, OH. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems. Suitable heteroatoms include oxygen, nitrogen, S(O)<sub>i</sub>, wherein i is 0, 1 or 2, and suitable heterocyclic aryls include furanyl, thienyl, pyridyl, pyrrolyl, pyrimidyl, pyrazinyl, imidazolyl, and the like.

"Heterocyclo" refers to a reduced heterocyclic ring system comprised of carbon, nitrogen, oxygen and/or sulfur atoms, and includes those heterocyclic systems described in "Handbook of Chemistry and Physics", 49th edition, 1968, R.C. Weast, editor; The Chemical Rubber Co., Cleveland, OH. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems.

"Heterocycloalkyl" refers to an alkyl group substituted with a heterocyclo group, and includes those heterocyclic systems described in "Handbook of Chemistry and Physics", 49th

edition, 1968, R.C. Weast, edit r; The Chemical Rubber Co., Cleveland, OH. See particularly Section C, Rules for Naming Organic Compounds, B. Fundamental Heterocyclic Systems. Preferably the alkyl group has from 1 to about 6 carbon atoms.

- 5 The term "lower" referred to herein in connection with organic radicals or groups defines such radicals or groups with one and up to and including 5 carbon atoms, preferably up to and including 4 carbon atoms, and advantageously one or two carbon atoms. Such radicals or groups may be straight  
10 chain or branched chain.


"Perfluoroalkyl" refers to an alkyl group which has every hydrogen replaced with fluorine.

"Perfluoroaryl" refers to an aryl group which has every hydrogen replaced with fluorine.


- 15 "Perfluoroaryl alkyl" refers an aralkyl group in which every hydrogen on the aryl moiety is replaced with fluorine.

"Pharmaceutically acceptable salt" includes salts of the compounds of the present invention derived from the combination of such compounds and an organic or inorganic  
20 acid. In practice the use of the salt form amounts to use of the base form. The compounds of the present invention are useful in both free base and salt form, with both forms being considered as being within the scope of the present invention.

- 25 The term "Arg-al" refers to the residue of L-argininal which has the formula:




The term "Arg-ol" refers to the residue of L-argininol which has the formula:




5

"(S)-N<sup>ω</sup>-nitroargininol hydrochloride" refers to the compound which has the formula:



10

"N-alpha-t-butoxycarbonyl-N<sup>ω</sup>-nitro-L-arginine" refers to the compound which has the formula:



15

"Bn" refers to benzyl.

"Boc" refers to t-butoxycarbonyl.

"BzlSO<sub>2</sub>" refers to benzylsulfonyl.

20

"Cbz" or "CBz" refers to benzyloxycarbonyl.

"DCA" refers to dichloroacetic acid.

"DCC" refers to N,N'-dicyclohexylcarbodiimide.

"DMF" refers to N, N-dimethylformamide.

- "DMSO" refers to dimethyl sulf oxide.  
"DMAP" refers to 4-N,N-dimethylaminopyridine.  
"EDC" refers to 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide hydrochloride salt.  
5 "Et<sub>3</sub>N" refers to triethylamine.  
"EtOH" refers to ethanol.  
"HBTU" refers to 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate.  
"HCl" refers to hydrochloric acid.  
10 "HOAc" refers to acetic acid.  
"HPLC" refers to high pressure liquid chromatography.  
"HOBT" refers to 1-hydroxybenzotriazole monohydrate.  
"i-BuOCOC1" refers to isobutylchloroformate.  
"LiAlH<sub>4</sub>" refers to lithium aluminum hydride.  
15 "LiAlH<sub>2</sub>(OEt)<sub>2</sub>" refers to lithium aluminum dihydride diethoxide.  
"Me" refers to methyl.  
"NMM" refers to N-methylmorpholine.  
"PhB(OH)<sub>2</sub>" refers to phenylboronic acid.  
20 "THF" refers to tetrahydrofuran.  
"TLC" refers to thin layer chromatography.

#### Brief Description of the Drawings

- 25 Figure 1 provides a synthetic scheme for compound 11, which is further detailed in Examples 1 to 11. In this figure, the steps are defined as follows: In the synthesis of (2) from 6-amino-m-cresol: 1. phthalic anhydride, acetic acid, reflux to give a 95% yield of (1), N-(2-hydroxy-4-methylphenyl)phthalimide); and 2. allyl bromide, K<sub>2</sub>CO<sub>3</sub>,  
30 acetonitrile, reflux to give 97% yield of (2). In the synthesis of (3) from (2): N,N-dimethylaniline, 185°C, about two hours, to give a 95% yield of (3). In the synthesis of (5) from (3): 1. benzylbromide, K<sub>2</sub>CO<sub>3</sub>, acetonitrile, reflux  
35 to give an 81% yield of (4), the benzyl ether derivative of (3); and 2. hydrazine, ethanol, reflux to give a 97% yield of

- (5). In the synthesis of (6) from (5):  $\alpha$ -toluene sulfonyl chloride, 4-methylmorpholine, acetonitrile, temperature 0°C, then warming to room temperature to give 99% yield of (6). In the synthesis of (8) from (6): 1. ozone gas, methanol, -78°C; 2. dimethylsulfide, -78°C, then warming to room temperature; 3. 1N HCl, acetone, room temperature then reflux to give an intermediate phenylacetaldehyde derivative (7); and 4. Jones' oxidation, acetone, 0°C, then warming to room temperature to give approximately quantitative yield of (8).
- 5 In the synthesis of (10) from (8): 1. hydrochloride salt of N<sup>9</sup>-nitro-L-argininal ethyl cyclol, EDC, 1-hydroxybenzotriazole hydrate, N,N-diisopropylethylamine, acetonitrile, to give a 69% yield of (9); and 2. H<sub>2</sub>, Pd/C catalyst, 55 PSI, ethanol, acetic acid, water to give a 89% yield of (10). In the synthesis of (11) from (10): 1. 5N HClO<sub>4</sub>, 0°C, 60 minutes; and 2. HPLC, to give a 62% yield of (11).
- 10  
15

- Figure 2 provides a synthetic scheme for compound 19, which is further detailed in Examples 12 to 14. In this figure, the steps are defined as follows. In the synthesis of (2) from 6-amino-M-cresol: 1. phthalic anhydride, acetic acid, reflux, to give a 95% yield of (1); and 2. allyl bromide, K<sub>2</sub>CO<sub>3</sub>, acetonitrile, reflux to give a 97% yield of (2). In the synthesis of (3) from (2). N,N-dimethylaniline, 185°C, about two hours, to give a 95% yield of (3). In the synthesis of (13) from (3): 1. iodomethane, K<sub>2</sub>CO<sub>3</sub>, acetonitrile, reflux, to give an 81% yield of (12); and 2. hydrazine, ethanol, reflux, to give a 90% yield of (13). In the synthesis of (14) from (13):  $\alpha$ -toluene sulfonyl chloride, 4-methylmorpholine, acetonitrile, temperature 0°C, then warming to room temperature, to give approximately quantitative yield of (14). In the synthesis of (16) from (14): 1. ozone gas, methanol, -78°C; 2. dimethylsulfide -78°C, then warming to room temperature, to give an intermediate phenylacetaldehyde (15); and 3. Jones' oxidation, acetone, 0°C, then warming to room temperature, to
- 20  
25  
30  
35

give a 46% overall yield of (16). In the synthesis of (18) from (16): 1. hydrochloride salt of N<sup>9</sup>-nitro-L-argininal ethyl cyclol, EDC, HOBT, N,N-diisopropylethylamine acetonitrile, to give a 62% yield of (17); and 2. H<sub>2</sub>, Pd/C catalyst, 55 PSI, ethanol, acetic acid, water, to give a 65% yield of (18). In the synthesis of (19) from (18): 1. 3N HCl, room temperature; and 2. HPLC, to give a 74% yield of (19).

Figure 3 depicts a reaction scheme for preparation of an intermediate used for the synthesis of certain compounds of formula (I). In this figure, "i" through "vi" are defined as: i) thionyl chloride, methanol; ii) di-tert-butyl dicarbonate, pH 7-8; iii) hydrogen gas, platinum oxide in ethanol, water and acetic acid; iv) bis-benzyloxycarbonyl S-methylisothiurea, base, tetrahydrofuran; v) calcium chloride, sodium borohydride in tetrahydrofuran and ethanol; vi) HCl (anhydrous). "\*" indicates the position of an asymmetric carbon atom.

Figures 4 and 5 depict preferred reaction schemes for synthesis of certain compounds of formula (I). In these figures, "i" through "iii" are defined as: i) HOBT, EDC, dimethylaminopyridine, triethylamine; ii) hydrogen gas, 10% palladium on charcoal and iii) dimethylsulfoxide, toluene, dichloroacetic acid and EDC.

Figure 6 depicts a reaction scheme for preparation of an intermediate used for the synthesis of certain compounds of formula (I). In this figure, "i" through "xvi" are defined as: (i) potassium iodide, dioxane; (ii) 2.5 M sodium ethoxide in ethanol, argon atmosphere; reflux 6 hours; (iii) yield after workup 60%; (iv) pyridine, triethylamine; (v) H<sub>2</sub>S(g), stirred at room temperature 16 hours; yield after workup 98%; (vi) acetone, iodomethane, reflux 30 minutes, filtration, methanol; (vii) ammonium acetate, reflux 1 hour, filter and dry; (viii) concentrated HCl, reflux 3 hours, yield after workup 30%; (ix) dioxane, sodium bicarbonate, di-

t-butyl dicarbonate, stir 18 hours at room temperature; (x) 4°C, 4.0 N NaOH to pH 12; 4-methoxy-2,3,6-trimethylbenzenesulfonyl chloride in dioxane; (xi) 1.0 N HCl to pH 7-8, water, yield after workup 68%; (xii) O,N-dimethylhydroxylamine hydrochloride, hydroxybenzotriazole hydrate, 4-methylmorpholine, THF, stir 2 hours, yield after workup 69%; (xiii)  $\text{LiAlH}_4$ , THF, dry ice/acetone bath; (xiv) aqueous potassium bisulfate, yield after workup 86%; (xv) 4-Benzhydrylsemicarbazide trifluoroacetate salt (the compound of Example 23), sodium acetate trihydrate in ethanol, reflux, yield after workup 89%; and (xvi) 50% TFA/DCM, add to ether, yield after workup 79%.

Figure 7 depicts a preferred reaction scheme for synthesis of certain compounds of formula (I). In this figure, "i" through "iii" are defined as: (i) EDC, hydroxybenzotriazole, 4-methylmorpholine in DMF; (ii) HF/anisole (9:1), aqueous acetic acid; and (iii) methanol, 1% HF(aqueous), formalin.

Figure 8 depicts a reaction scheme for the preparation of a compound of formula (II) which incorporates an arginine chloromethylketone moiety at P1. In this figure, "i" through "iii" are defined as: (i) EDC, HOBT, NMM,  $\text{CH}_3\text{CN}$ , room temperature; (iia)  $i\text{-BuOCOC1}$ , NMM, DMF at  $-20^\circ\text{C}$ ; (iib) add 45, DMF,  $\text{Et}_3\text{N}$ , allow temperature to increase from  $-20^\circ$  to room temperature; and (iii) HF, anisole.

Figure 9 depicts a reaction scheme for the preparation of a compound of formula (II) which incorporates an arginine ketoamide moiety at P1. In this figure, "i" through "iii" are defined as: (i) EDC, HOBT, NMM,  $\text{CH}_3\text{CN}$ , room temperature; (ii)  $\text{H}_2$ , palladium on carbon, EtOH,  $\text{H}_2\text{O}$ , HOAc to give intermediate 50; and (iii) EDC, DMSO, DCAA, toluene, temperature is allowed to increase from  $0^\circ$  to room temperature.

Figure 10 depicts a reaction scheme for the preparation of a compound of formula (II) having a boronic acid moiety at




- P1. In this figure, "i" through "vii" are defined as: (i) treatment of 16 with  $i\text{BuCOCl}$ , NMM, THF, at  $-20^{\circ}\text{C}$ ; (ii) addition of 52,  $\text{CHCl}_3$ ; (iii)  $\text{Et}_3\text{N}$ , at  $-20^{\circ}\text{C}$ ; (iv)  $\text{NaN}_3$ , DMF, at temperature of 80 to  $100^{\circ}\text{C}$  to give intermediate azide 54;  
5 (v)  $\text{H}_2$ ,  $\text{Pd}(\text{OH}_2)/\text{carbon}$ , methanol,  $\text{HCl}$ ; (vi) formamidine sulfonic acid ( $\text{H}_2\text{NCl}=\text{NH}$ )  $\text{SO}_3\text{H}$ , DMAP,  $\text{EtOH}$ , reflux, to give intermediate 56; and (vii)  $\text{PhB}(\text{OH})_2$ ,  $\text{Et}_2\text{O}$ ,  $\text{H}_2\text{O}$ .

- Figure 11 depicts a reaction scheme for the preparation of a compound of formula (II) having a  $\text{C}_1$  to  $\text{C}_4$  perfluoroalkylargininyl ketone moiety at P1. In this figure,  
10 "i" to "vi" are defined as: (i) EDC, HOBT, NMM,  $\text{CH}_3\text{CN}$ , at room temperature; (ii)  $i\text{-BuOCOCl}$ , NMM, DMF, temperature about  $-30^{\circ}\text{C}$  to  $-15^{\circ}\text{C}$  to room temperature; (iii) TFA,  $\text{CH}_2\text{Cl}_2$ ,  
15 temperature about  $0^{\circ}\text{C}$  to room temperature to give intermediate 60; (iv) bis-Cbz-S-methyl isothioureia ( $\text{CbzNHC}(=\text{NCbz})\text{SMe}$ ), THF,  $\text{Et}_3\text{N}$ ; (v) Swern oxidation with mixture of oxalylchloride, DMSO,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , temperature at about  $-78^{\circ}\text{C}$  to room temperature, to give ketone intermediate 62; and (vi) HF, anisole, temperature at about  $-78^{\circ}\text{C}$  to room temperature.

- 20 Figure 12 depicts the reaction scheme for preparation of intermediate 69 which is used for the synthesis of certain compounds of formula (I). In this figure "i" through "vi" are defined as: i) thionyl chloride, methanol; ii) di-tert-butyl carbonate, pH 7-8; iii) hydrogen gas, platinum oxide in  
25 ethanol, water and acetic acid; iv) bis-benzyloxycarbonyl 5-methylisothioureia, base, tetrahydrofuran; v) calcium chloride, sodium borohydride in tetrahydrofuran and ethanol; vi)  $\text{HCl}$  (anhydrous). "\*" indicates the position of an assymetric carbon atom.

Detailed Description of the Invention1. Preferred Novel Compounds

Novel compounds of the present invention include  
 5 compounds of the Formula (I):



wherein

10 (a) X is selected from the group consisting of  $-S(O)_2-$ ,  $-N(R')-S(O)_2-$ ,  $-C(=O)-$ ,  $-OC(=O)-$ ,  $-NH-C(=O)-$ ,  $-P(O)(R'')$  and a direct link, wherein R' is hydrogen, alkyl of 1 to about 4 carbon atoms, aryl of about 6 to about 14 carbon atoms or aralkyl of about 6 to about 16 carbon atoms, and R'' is NHR',  
 15 OR', R', or SR';

(b) R<sub>1</sub> is selected from the group consisting of:

(1) alkyl of 1 to about 12 carbon atoms,  
 (2) alkyl of 1 to about 3 carbon atoms substituted  
 with cycloalkyl of about 5 to about 8 carbon atoms, which  
 20 optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,

(3) cycloalkyl of 3 to about 15 carbon atoms, which optionally is substituted on the ring carbons with  
 25 hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,

(4) heterocycloalkyl of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group  
 30 consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, wherein i is 0, 1

or 2, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

5 (5) heterocyclo of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub> and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl or alkoxy, or -CO<sub>2</sub>R',

10 (6) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 5 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

15 (7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,

(8) heteroaryl of 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,





25 (9) aralkyl of about 7 to about 15 carbon atoms having 6 to about 14 ring atoms which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

30 (10) heteroaralkyl of 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

35 (11) aralkenyl of about 8 to about 16 carbon atoms having 6 to about 14 ring atoms which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or

Y<sub>3</sub>,

- (12) heteroaralkenyl of 5 to about 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,



- (17) perfluoroalkyl of 1 to about 12 carbon atoms,

- (18) perfluoroaralkyl of about 6 to about 14 carbon atoms,

- (19) perfluoroaralkyl of about 7 to about 15 carbon atoms,

- (20) hydrogen, and

- (21) , wherein is a 5 to 7 membered heterocyclic ring having 3 to 6 ring carbon atoms, where V is -CH<sub>2</sub>-, -O-, -S(=O)-, -S(O)<sub>2</sub>- or -S-, and


wherein  $Y_1$ ,  $Y_2$ , and  $Y_3$  are

- (i) independently selected from the group consisting of hydrogen, halogen, cyano, nitro, tetrazolyl, amino, guanidino, amidino, methylamino, methylguanidino,   
 5 -CF<sub>3</sub>, -CF<sub>2</sub>CF<sub>3</sub>, -CF<sub>2</sub>H, -CH(CF<sub>3</sub>)<sub>2</sub>, -C(OH)(CF<sub>3</sub>)<sub>2</sub>, -OCF<sub>3</sub>,  
 -OCF<sub>2</sub>CF<sub>3</sub>, -OC(O)NH<sub>2</sub>, -OC(O)NHZ<sub>1</sub>, -OC(O)NZ<sub>1</sub>Z<sub>2</sub>, -NHC(O)Z<sub>1</sub>,  
 -NHC(O)NH<sub>2</sub>, -NHC(O)NHZ<sub>1</sub>, -NHC(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OH, C(O)NH<sub>2</sub>,  
 -C(O)NHZ<sub>1</sub>, -C(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OZ<sub>1</sub>, -P(O)<sub>3</sub>H, -P(O)<sub>3</sub>H<sub>2</sub>, -PH(O)OH,  
 -P(O)<sub>3</sub>(Z<sub>1</sub>)<sub>2</sub>, -S(O)<sub>3</sub>H, -S(O)<sub>m</sub>Z<sub>1</sub>, -Z<sub>1</sub>, -OZ<sub>1</sub>, -OH, -NH<sub>2</sub>, -NHZ<sub>1</sub>,  
 10 and -NZ<sub>1</sub>Z<sub>2</sub>, wherein m is 0, 1 or 2, and Z<sub>1</sub> and Z<sub>2</sub> are  
 independently selected from the group consisting of alkyl of  
 1 to about 12 carbon atoms, aryl of about 6 to about 14  
 carbon atoms, heteroaryl of about 5 to about 14 atoms having  
 1 to about 9 carbon atoms, aralkyl of about 7 to about 15  
 15 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms  
 having about 3 to about 9 carbon atoms, or

- (ii)  $Y_1$  and  $Y_2$  are selected together to be  
 -O[C(Z<sub>3</sub>)(Z<sub>4</sub>)]<sub>q</sub>O-, wherein q is an integer from 1 to 4 and Z<sub>3</sub>  
 and Z<sub>4</sub> are independently selected from the group consisting  
 20 of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of  
 about 6 to about 14 carbon atoms heteroaryl of about 5 to  
 about 14 atoms having 1 to about 9 carbon atoms, aralkyl of  
 about 7 to about 15 carbon atoms, and heteroaralkyl of about  
 6 to about 11 atoms having about 3 to about 9 carbon atoms;

- (c) R<sub>2</sub> is selected from the group consisting of  
 25 hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of  
 about 2 to about 4 carbon atoms;

(d) R<sub>3</sub> is selected from the group consisting of



- 30 where d is an integer from 0 to 5 and W is -N- or -CH-;

(e) R<sub>4</sub> is selected from the group consisting of

- (i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein n is 0, 1

or 2; and  $R_1$  is as defined above, with the proviso that  $R_1$  is not a camphor derivative or  $\text{---}\overset{\text{N}}{\text{V}}$  heterocycle,

(ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ ,  $-\text{OCF}_2\text{H}$ ,

(iii) halogen,

5 (iv) alkyl of 1 to 6 carbon atoms substituted with  $Z_5$ , wherein  $Z_5$  is selected from the group consisting of hydrogen,  $-\text{OH}$ ,  $-\text{OR}_8$ ,  $-\text{CH}_3$ , halogen,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{OR}_8$  and  $-\text{S}(\text{O})_p\text{R}_8$ , wherein  $R_8$  is alkyl of 1 to about 6 carbon atoms, and  $p$  is 0, 1 or 2,

10 (v) alkyl of 1 to 3 carbon atoms substituted with cycloalkyl of 3 to 5 carbon atoms,

(vi) alkenyl of about 3 to about 6 carbon atoms,

(vii) cycloalkyl of about 3 to about 10 carbon  
atoms,

15 (viii) heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms are heteroatoms selected from oxygen, sulfur and nitrogen, and

(ix) heteroaralkyl of about 5 to about 10 ring  
20 atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen;

(f)  $R_5$  is selected from the group consisting of

(i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ ,  $R_1$  is as defined  
25 above, with the proviso that  $R_1$  is not a camphor derivative

or  $\text{---}\overset{\text{N}}{\text{V}}$  heterocycle,

(ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ , and

(iii) halogen;

(g)  $R_6$  is selected from the group consisting of

30 (i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein  $R_1$  is as defined above, with the proviso that  $R_1$  is not a camphor

derivative or  $\text{---}\overset{\text{N}}{\text{V}}$  heterocycle,

(ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ ,

(iii) halogen,

(iv) alkyl of 1 to about 12 carbon atoms substituted with  $Z_6$ , wherein  $Z_6$  is selected from the group consisting of hydrogen, hydroxy, halogen,  $-OR_9$ ,  $-NHR_9$ ,  $-C(O)OH$ ,  $-C(O)OR_9$ , and  $-S(O)_pR_9$ , wherein  $R_9$  is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon atoms, aralkyl of about 7 to about 12 carbon atoms, heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ ; or heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms including 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, and which is optionally mono-, di- or tri- substituted on the ring carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ , and  $p$  is 0, 1 or 2;

(v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms;

(vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 6 to about 10 carbon atoms or heteroaryl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms,

(vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri- substituted on the ring carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ ,

(viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri- substituted on the ring carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ ,

(ix) aralkenyl of 6 to about 15 carbon atoms which

is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

(x) heteroaralkenyl of about 5 to about 10 ring atoms having 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>; and

(h) R<sub>7</sub> is selected from the group consisting of

(i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above with the

proviso that R<sub>1</sub> is not a camphor derivative or heterocycle,

(ii) -CF<sub>3</sub>, or -CF<sub>2</sub>H,

(iii) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>,

(iv) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(vi) heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, and

(vii) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>, and wherein each R<sub>10</sub> and R<sub>11</sub> is independently selected and is alkyl of 1 to about 4 carbon atoms or hydrogen; or pharmaceutically acceptable salts thereof.

Preferred X groups include a direct link, -SO<sub>2</sub>-, and -NH-S(O)<sub>2</sub>-. An especially preferred X group is -SO<sub>2</sub>-.

Preferred R<sub>1</sub> groups include alkyl, aralkyl, and aryl




groups. Preferred aryl groups include substituted or unsubstituted phenyl and naphthyl, 2,3-methylenedioxyphenyl, and 3,4-methylenedioxyphenyl. Preferred substitutions include, methyl, methoxy, fluoro, chloro, trifluoromethyl, and -OCF<sub>3</sub>. Meta and ortho ring substitutions are preferred. Meta and ortho di-substitutions on the ring are also preferred.

Particularly preferred R<sub>1</sub> groups include aralkyl groups. Especially preferred R<sub>1</sub> groups include substituted or unsubstituted benzyl and naphthylalkyl groups. Cyclohexyl and cyclohexylmethyl are also especially preferred R<sub>1</sub> groups.

A particularly preferred R<sub>2</sub> group is hydrogen.

Preferred R<sub>3</sub> groups include



Preferred R<sub>4</sub> groups include:

- (i) hydrogen,
- (ii) alkyl of 1 to 6 carbon atoms substituted with Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of hydrogen, -OH, -OR<sub>8</sub>, -CH<sub>3</sub>, halogen, -C(O)OH, -C(O)OR<sub>8</sub> and -S(O)<sub>p</sub>R<sub>8</sub>;
- (iii) alkyl of 1 to 3 carbon atoms substituted with cyclic alkyl of 3 to 5 carbon atoms,
- (iv) alkenyl of about 3 to about 6 carbon atoms,
- (v) cycloalkyl of about 3 to about 7 carbon atoms,
- (vi) heteroaryl having 1 to about 4 ring carbon atoms and the remainder of the ring atoms are heteroatoms selected from oxygen, sulfur and nitrogen,
- (vii) heteroaralkyl of 2 to about 6 carbon atoms having 1 to about 4 ring carbon atoms and the remainder of the ring atoms are heteroatoms selected from oxygen, sulfur and nitrogen, and

(viii) trifluoromethyl.

Preferred R<sub>4</sub> groups also include methyl, ethyl, propenyl, allyl, propyl, isopropyl, butyl, R-sec-butyl, S-sec-butyl, isobutyl, 1-pentyl, R-2-pentyl, S-2-pentyl, 5 3-pentyl, S-1-(2-methyl)-butyl, R-2-(3-methyl)-butyl, 1-(3-methyl)-butyl, R-1-(2-methyl)-butyl, cyclopentyl, 2-pyrrolyl, 3-pyrrolyl, 1-hexyl, S-2-hexyl, R-2-hexyl, R-3-hexyl, S-3-hexyl, hydroxy, lower alkoxy of 1 to about 4 carbon atoms, phenyloxy, -OCF<sub>3</sub>, and halogen.

10 Preferred R<sub>5</sub> groups include hydrogen, halogen, alkyl of 1 to about 5 carbon atoms, trifluoromethyl, and alkoxy of 1 to about 4 carbon atoms. Hydrogen is an especially preferred R<sub>5</sub> group.

Preferred R<sub>6</sub> groups include:

15 (i) hydrogen,

(ii) alkyl of 1 to about 12 carbon atoms substituted with Z<sub>6</sub>, wherein Z<sub>6</sub> is as defined above,

(iii) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 5 to about 8 carbon 20 atoms,

(iv) alkenyl of about 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 6 to about 10 carbon atoms or heteroaryl of 1 to about 9 carbon atoms,

25 (v) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(vi) heteroaralkyl of about 5 to about 10 ring 30 atoms having 1 to about 9 ring carbon atoms and having the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or 35 Y<sub>3</sub>,

(vii) aralkenyl of about 6 to 15 carbon atoms

which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

- (viii) heteroaralkenyl of about 5 to about 10 ring atoms having 1 to 9 ring carbon atoms and having the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>.

More preferred R<sub>6</sub> groups, when R<sub>4</sub> and R<sub>5</sub> are independently hydrogen or methyl, are selected from the group consisting of aralkyl of about 8 to about 13 carbon atoms, -O-R<sub>1</sub>, -NH-R<sub>1</sub>, and -S(O)<sub>p</sub>R<sub>1</sub> wherein R<sub>1</sub> is preferably aralkyl of about 7 to about 12 carbon atoms. Preferred aryl components of the aralkyl groups include unsubstituted and substituted phenyl or naphthyl groups. Preferred substitutions on the aryl ring include methyl, methoxy, fluoro, chloro, trifluoromethyl, alkoxy carbonyl, methylenedioxy and amidocarbonyl. Phenylethyl, phenylpropyl, cyclohexylethyl and cyclohexylpropyl are especially preferred R<sub>6</sub> groups.

Especially preferred R<sub>7</sub> groups include hydrogen, methyl, difluoromethyl and trifluoromethyl. Hydrogen and methyl are especially preferred R<sub>7</sub> groups. Preferred R<sub>7</sub> groups also include those selected from the group consisting of:

(i) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(ii) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,


(iii) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(iv) aryl and heteroaryl of about 5 to about 10 ring atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, and

(v) alkyl of 1 to about 3 carbon atoms substituted with heter aryl of ab ut 5 t about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>.

5 A particularly preferred embodiment is when R<sub>5</sub> and R<sub>6</sub> are independently selected to be hydrogen or methyl, and R<sub>4</sub> is selected from the group consisting of hydrogen, methyl, ethyl, propenyl, allyl, propyl, isopropyl, butyl, R-sec-butyl, S-sec-butyl, isobutyl, 1-pentyl, R-2-pentyl, S-2-pentyl, 3-pentyl, S-1-(2-methyl)-butyl, R-2-(3-methyl)-butyl, 10 1-(3-methyl)-butyl, R-1-(2-methyl)-butyl, cyclopentyl, 2-pyrrolyl, 3-pyrrolyl, 1-hexyl, S-2-hexyl, R-2-hexyl, R-3-hexyl, and S-3-hexyl. A particularly preferred embodiment according to this aspect includes compounds where R<sub>4</sub> is 15 hydrogen or methyl.

According to a particularly preferred aspect, provided are compounds of formula I wherein X is -S(O)<sub>2</sub>-, R<sub>1</sub> is substituted or unsubstituted aralkyl, and R<sub>3</sub> is .



A very preferred aspect is directed to such compounds where R<sub>1</sub> is substituted or unsubstituted benzyl.


25 According to another preferred aspect, provided are compounds of formula I, wherein any two of R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, and R<sub>6</sub> are hydrogen.

Particularly preferred compounds according to this invention include N-[2-hydroxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-L-argininal (Example 11) and N-[2-methoxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-L-argininal (Example 14).

30 According to another aspect, the present invention is directed to compounds having inhibitory activity toward

pr teases of the trypsin/chymotrypsin class. Crystal structures of serine protease inhibitors in complex with their cognate enzymes have been determined and are available from the Brookhaven National Laboratory Protein Data Bank.

- 5 The three-dimensional crystal structures of proteases of the trypsin/chymotrypsin class are highly conserved within the class. The most potent inhibitors in the class make specific hydrogen bonds with structurally conserved backbone atoms of amino acid 216 of the protease (using the standard  
10 chymotrypsin numbering system). In most inhibitors, these hydrogen bonds are formed by atoms of peptide amide bonds or sulfonamide bonds. Thus, this aspect of the invention is directed to compounds of formula (II):



- 15 wherein a substituted phenol or phenyl ether is present in the compound in place of the amide bond of an inhibitor that makes hydrogen bonds to amino acid 216 in the trypsin/chymotrypsin class of serine proteases. Thus, according to this aspect of the present invention, provided  
20 are compounds of formula (II), wherein X, R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are as defined hereinabove in connection with formula (I), and

- (a) R<sub>12</sub> is selected from the group consisting of
- (i) alkyl of 1 to about 12 carbon atoms optionally substituted with amino, amidino, or guanidino,
  - (ii) aralkyl of 7 to about 15 carbon atoms which is optionally substituted on a ring carbon with amino, amidino or guanidino,
  - (iii) aryl of 6 to about 14 carbon atoms which is
- 30 optionally substituted on a ring carbon with amino, amidino,

or guanidino, and

(iv) heteroaralkyl of 5 to about 14 ring atoms having about 1 to about 13 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur which is optionally substituted on a ring carbon with amino, amidino or guanidino; and

(b) TS is selected from the group consisting of

- (i) -CHO,
- (ii) -C(O)CF<sub>3</sub>,
- (iii) -C(O)CF<sub>2</sub>CF<sub>3</sub>,
- (iv) -C(O)-C(O)-NHR<sub>13</sub>,
- (v) -C(O)-C(O)<sub>2</sub>R<sub>13</sub>,
- (vi) -C(O)-C(O)-NR<sub>13</sub>R<sub>14</sub>,
- (vii) -B(OR<sub>13</sub>)(OR<sub>14</sub>),
- (viii) -C(O)CH<sub>2</sub>Cl, and
- (ix) -C(O)R<sub>15</sub>;

wherein R<sub>13</sub> and R<sub>14</sub> are independently selected from hydrogen lower alkyl of 1 to about 6 carbon atoms, cycloalkyl of 3 to about 10 carbon atoms, aralkyl of 7 to about 15 carbon atoms, and heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur, and R<sub>15</sub> is selected from aryl of 6 to about 14 carbon atoms and heteroaryl of 5 to about 14 ring atoms having 1 to about 13 ring carbon atoms and the remainder of the ring atoms selected from oxygen, nitrogen and sulfur. In the case where TS is (vii), additionally -OR<sub>13</sub> and -OR<sub>14</sub> may be taken together to be -O[C(Z<sub>3</sub>)(Z<sub>4</sub>)]qO- wherein Z<sub>3</sub>, Z<sub>4</sub> and q are as defined in connection with formula (I). Also included are pharmaceutically acceptable salts of the compounds of formula (II).

According to a preferred aspect, inhibitors of elastase

will have a small alkyl group, such as methyl, for  $R_{12}$ .

Inhibitors of chymotrypsin preferably will have an aralkyl group, such as benzyl, for  $R_{12}$  according to formula (II).

Serine protease inhibitors that have a basic amino acid in the primary specificity site, such as inhibitors of trypsin, thrombin, factor Xa, factor VIIa, will have an amino, amidino, or guanidino substituted alkyl, aryl, or aralkyl group for  $R_{12}$ .

According to another aspect, the present invention includes within its scope salts of the compounds of formulas I and II. These salts include salts of the compounds of the present invention derived from the combination of such compounds and an organic or inorganic acid. In practice, the use of the salt form may amount to use of the (free) base form. The compounds of the present invention are useful in both free base and salt form, with both forms being considered as being within the scope of the present invention. These salts include acid addition salts, for example, salts of hydrochloric acid, hydrobromic acid, trifluoroacetic acid, acetic acid, benzene sulfonic acid and other suitable acid addition salts including maleate, fumarate and citrate salts.

## 2. Preparation of Preferred Compounds

Figures 1 and 2 depict synthetic schemes for two preferred compounds according to this invention. Figure 1 outlines a sequence of reactions for the preparation of a preferred species of the present invention. Commercially available 6-amino-m-cresol is reacted with phthalic anhydride, optionally in the presence of an inert solvent like acetic acid, toluene, or xylene, at about 100-150°C to afford the phthalimide intermediate 1. Reaction of 1 with allyl bromide in the presence of a suitable base such as anhydrous potassium carbonate in an inert solvent such as N,N-dimethylformamide, dioxane, tetrahydrofuran, or

acetonitrile at a temperature in the range from about room temperature to about 110°C provides the allyl ether derivative intermediate 2. Thermal Claisen rearrangement of 2, performed either neat or optionally in a suitable inert solvent such as N,N-dimethylaniline, at about 150-250°C gives the allylphenol 3. Reaction of phenol derivative 3 with benzyl bromide in an inert solvent such as N, N-dimethylformamide, dioxane, tetrahydrofuran, or acetonitrile at about room temperature to about 110°C provides the benzyl ether derivative 4. Cleavage of the phthalimido moiety of 4 is achieved selectively utilizing hydrazine in an alcoholic solvent such as ethanol at a temperature from about 20 to 80°C and affords the aniline derivative 5. Coupling of 5 with a suitable sulfonyl chloride such as benzylsulfonyl chloride in an inert solvent such as N, N-dimethylformamide, dioxane, tetrahydrofuran, or acetonitrile at a temperature of about -10°C to about 50°C in the presence of a base such as pyridine, collidine, triethylamine, N,N-diisopropylethylamine, or N-methylmorpholine gives the benzylsulfonamide derivative 6. Treatment of a solution of 6 in solvent, preferably methanol, ethanol, ethyl acetate, acetone or methylene chloride, or mixtures of the preceding solvents, with a stream of ozone gas at a temperature of about -80°C to about 0°C, followed by reductive decomposition of the ozonide *in situ* with reagents such as dimethyl sulfide, triethyl phosphite, triphenylphosphine, triethylamine, or zinc metal generates an aldehyde intermediate 7. When alcoholic solvents such as methanol are utilized, the reaction affords a mixture of the desired aldehyde along with varying amounts of the corresponding dimethyl acetal derivative. These mixtures are dissolved in a inert solvent such as acetone and hydrolyzed with dilute mineral acids such as 1 N hydrochloric acid at about room temperature to about reflux temperature which quantitatively affords the desired aldehyde 7. Oxidation of aldehyde substrate 7 with reagents such as pyridinium dichromate in



N,N-dimethylformamide or chromic acid in acetone at a temperature of about 0°C to 30°C gives the carboxylic acid derivative 8. Compound 8 is coupled to N<sup>ε</sup>-nitro-L-argininal ethyl cyclol using 1-hydroxybenzotriazole monohydrate, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride salt, and then N,N-diisopropylamine to give 9. The N<sup>ε</sup>-nitro group of 9 is removed by treatment with hydrogen gas and palladium on carbon in ethanol, water, and acetic acid at about 20 to 60 psi. This reaction produces acetate acid salt 10. The ethyl cyclol group of 10 is hydrolyzed by treatment with a suitable acid catalyst such as dilute hydrochloric acid, sulfuric acid, methanesulfonic acid, perchloric acid or hexafluorophosphoric acid followed by HPLC purification with 0.1% trifluoroacetic acid in acetonitrile and water to produce the product 11.

The reaction scheme of Figure 2 depicts the preparation of another very preferred compound. It differs from the synthetic scheme of Figure 1 in that intermediate 3 is first reacted with methyl iodide in the presence of a suitable base such as anhydrous potassium carbonate in an inert solvent such as N,N-dimethylformamide, dioxane, tetrahydrofuran, or acetonitrile at a temperature of about room temperature to about 110°C, to provide the methyl ether derivative 12. Intermediate 12 is processed through an analogous set of reactions to those outlined in Figure 1 to produce the target argininal derivative 19.

The preferred means of chemically coupling (as for example, intermediate 8 to argininal ethyl cyclol to give intermediate 9 as depicted in Figure 1) include formation of a peptide bond using conventional coupling reagents known in the art. See Bodanszky, N., *Peptide Chemistry*, pp. 55-73, Springer-Verlag, New York (1988) and references cited therein. The chemical coupling may be either by means of one-step or two-step coupling. In one-step coupling, the two coupling partners are coupled directly. Preferred coupling reagents for one-step coupling include DCC with HOBt, EDC

with HOBt, BOP, HBTU or TBTU. In two-step coupling, an activated ester or anhydride of the C-terminal carboxy group of one coupling partner is formed prior to its coupling to the other coupling partner.

5 For compounds of the present invention containing alkenyl or aryl moieties substituted with halogen, cyano, nitro, or -S-Z<sub>1</sub>, it is preferred to avoid the use of hydrogen gas with palladium on carbon for cleaving the N<sup>9</sup>-nitro protecting group on the arginine side chain. Instead, it is  
10 preferred to use boron tris(trifluoroacetate), B(OCOCF<sub>3</sub>)<sub>3</sub>, to cleave the N<sup>9</sup>-nitro of the arginine group. This reagent may be prepared by the reaction of BBr<sub>3</sub> and CF<sub>3</sub>COOH in dichloromethane at 0°C. Alternatively, the reagent is also  
15 commercially available. Generally, the N<sup>9</sup>-nitro compound is treated with boron tris(trifluoroacetate) in trifluoroacetic acid at 0°C. See, e.g., Fieser, M. and Fieser, L. F., Reagents for Organic Synthesis, p. 46, John Wiley & Sons, New York (1974); and Pless, J., and Bauer, W. Angew. Chem., Internat. Ed., 12, 147 (1973).

20 Another preferred reagent for selective nitro group cleavage is titanium trichloride. This reagent is commercially available. The N<sup>9</sup>-nitro compound is treated with titanium trichloride in aqueous methanol containing an ammonium acetate buffer followed by exposure of the reaction  
25 mixture to air or dimethyl sulfoxide. See, e.g., Freidinger, R.M., Hirschmann, R., and Veber, D.F., J. Org. Chem., 43, 4800 (1978).

Another preferred method for synthesizing these compounds having an L-argininal moiety is to use the di-N-t-  
30 butoxycarbonyl protecting group for the L-argininal moiety for groups incompatible with hydrogenation with palladium on carbon. For example, alpha-N-benzyloxycarbonyl-omega, omega'-di-N-t-butoxycarbonylarginine is dissolved in acetonitrile and treated with hydroxybenzotriazole and 1-  
35 ethyl-3-(3-dimethylamino-propyl)carbodiimide HCl salt to form alpha-N-benzyloxycarbonyl-omega, omega'-di-N-t-

butoxycarbonyl-L-arginine lactam. The lactam is reduced by treatment with  $\text{LiAlH}_4$  in THF at  $-70^\circ\text{C}$  to provide alpha-N-benzyloxycarbonyl-omega, omega'-di-N-t-butoxycarbonyl-L-argininal. This aldehyde is protected as the diethyl acetal by treatment with ethanol and HCl. The N-benzyloxycarbonyl protecting group is removed by treatment with hydrogen gas and palladium on carbon to give omega, omega'-di-N-t-butoxycarbonyl-L-argininal diethyl acetal, HCl salt. This protected L-argininal moiety can then be coupled to a desired carboxylic acid by treatment with N-hydroxybenzotriazole and 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide HCl salt. The diethyl acetal and the di-Boc protecting groups are removed by treatment with hexafluorophosphoric acid in acetonitrile at  $0^\circ\text{C}$ . The reaction mixture is quenched by addition of 2.5 M aqueous sodium acetate until pH 4 is reached. The mixture is filtered through a 2 micron filter. Preparative HPLC using 0.1%  $\text{CF}_3\text{COOH}$  in 10-40% aqueous acetonitrile provides the trifluoroacetate salt of the desired substituted L-argininal compound.

Figure 3 depicts a reaction scheme for the preparation of intermediate 25 which is used in the synthesis of compounds of formula (I) according to reaction schemes depicted in Figures 4 and 5. Examples 15 to 19 describe preparation of 25 in further detail.

Figures 4 and 5 depict reaction schemes for the preparation of certain compounds of formula (I) wherein  $\text{R}_3$  is a 3-piperidinyl-(N-guanidino) group. Preparation of compound 28 is described in further detail in Examples 20 to 22. Compound 31 is prepared using the appropriate reagents according to procedures similar to those described in Examples 20 to 22.

Figure 6 depicts a reaction scheme for preparation of intermediate 41 which is used in the synthesis of compounds of formula (I) according to the reaction scheme depicted in Figure 7. Examples 24 to 30 describe the preparation of 41

in further detail.

Intermediates used in the preparation of compounds of formula (I) having a 4-amidinophenylalaninal group at P1 may be prepared according to the reaction scheme depicted in Figure 6 and described in Examples 24 to 30, using the appropriate alpha-bromo-para-tolunitrile starting material.

Figure 7 depicts a reaction scheme for preparation of a compound of formula (I) wherein R<sub>3</sub> is 3-amidinophenyl. Preparation of 44 is more fully described in Examples 31 to 33.

Figure 12 depicts a reaction scheme for the preparation of intermediate 69 which is used in the synthesis of compounds of formula (I). Intermediate 69 is prepared by procedures similar to those described in Examples 15 to 19 and using the appropriate starting materials. Intermediate 69 may be coupled to intermediates such as 8 and 16 according to reaction schemes analogous to those depicted in Figures 4 and 5.

Compounds of formula (II) may be prepared by methods such as described hereinbelow:

Figure 8 outlines a synthetic sequence for the preparation of a preferred compound of the invention which incorporates a P1-arginine chloromethylketone moiety. Coupling of carboxylic acid derivative 16 with the known arginine chloromethylketone precursor 45 (Kettner, C. and Shaw, E., *Biochemistry*, 17: 4778 (1978); Kettner, C. and Shaw, E., *Biochim. Biophys. Acta*, 569: 31 (1979)) under standard carbodiimide conditions of EDC, HOBT, and NMM at about 0°C to room temperature or under standard mixed anhydride conditions of isobutyl chloroformate and NMM at about -30°C to room temperature affords intermediate 46. Suitable solvents for either reaction include THF, acetonitrile or DMF. Double deprotection of intermediate 46 with anhydrous hydrogen fluoride optionally in the presence

of anisole at a temperature in the range of about -78°C to about room temperature gives the desired product 47.

Figure 9 outlines a synthetic sequence for the preparation of a preferred compound of the invention which incorporates a P1-ketoargininamide moiety. Coupling of carboxylic acid derivative 16 with the known hydroxyargininamide precursor 48 (see, e.g. commonly assigned U.S. Patent Nos. 5,371,072 and 5,597,804) under standard carbodiimide conditions of EDC, HOBT, and NMM at about 0°C to room temperature or under standard mixed anhydride conditions of isobutyl chloroformate, and NMM at about -30°C to room temperature affords intermediate 49. Suitable solvents for either reaction include THF, acetonitrile or DMF. Deprotection of intermediate 49 with anhydrous hydrogen fluoride optionally in the presence of anisole at about -78°C to about room temperature, or by catalytic hydrogenation using a suitable catalyst such as Pd/C or Pd(OH)<sub>2</sub>/C in a solvent composed of ethanol, water and/or acetic acid gives the intermediate 50. Finally, oxidation of intermediate 50 under the standard Moffatt conditions with EDC, DMSO and dichloroacetic acid in a suitable solvent such as toluene at about -10°C to room temperature produces the desired product 51.

Figure 10 outlines a synthetic sequence for the preparation of a preferred compound of the invention which incorporates a P1-boronic acid moiety. Coupling of carboxylic acid derivative 16 with the known protected boroarginine ester precursor 52 (See, e.g., Wityak, J. et. al., J. Org. Chem., 60: 3717 (1995) and references cited therein) under standard carbodiimide conditions of EDC, HOBT, and NMM at about 0°C to room temperature or under standard mixed anhydride conditions of isobutyl chloroformate, NMM and then *in situ* added Et<sub>3</sub>N at about -30°C to room temperature affords intermediate 53. Suitable solvents for either reaction include THF, acetonitrile, dichloromethane,

chl reform, DMF, or mixtures thereof. Reaction f  
intermediate 53 with sodium azide in a suitable inert solvent  
such as DMF, N, N-dimethylacetamide, DMSO or 1-methyl-2-  
pyrrolidinone at about room temperature to about 100°C affords  
5 the azide 54. Catalytic hydrogenation of intermediate azide  
54 in an alcoholic solvent such as methanol preferably in the  
presence of hydrogen chloride gives the intermediate amine  
salt 55. Reaction of intermediate 55 with formamidine  
sulfonic acid in an alcoholic solvent such as ethanol  
10 preferably in the presence of a tertiary amine, especially  
DMAP, at about room temperature to 100°C produces intermediate  
56. Deprotection of intermediate 56 by an exchange reaction  
preferably with phenylboronic acid in a two-phase solvent  
system such as ether and water then gives the desired boronic  
15 acid product 57.

Figure 11 outlines a synthetic sequence for the  
preparation of a preferred compound of the invention which  
incorporates a P1-C1 to C4 perfluoroalkylargininyl ketone  
moiety. Coupling of carboxylic acid derivative 16 with the  
20 known perfluoroargininol precursor 58 (see, e.g., Schacht,  
A.L., Shuman, R.T., and Wiley, M.R., UK Patent Application GB  
2287027 A, Sept. 6, 1995 and references cited therein; for  
the application of the Dakin-West protocol to arginine and  
related substrates, see also Ueda, T., Kam, C.M., and Powers,  
25 J.C., Biochem. J., 265: 539 (1990); Buchanan, G.L., Chem.  
Soc. Rev., 17: 91 (1988)) under standard carbodiimide  
conditions of EDC, HOBT, and NMM at about 0°C to room  
temperature or under standard mixed anhydride conditions of  
isobutyl chloroformate, and NMM at about -30°C to room  
30 temperature affords intermediate 59. Suitable solvents for  
either reaction include THF, acetonitrile, dichloromethane,  
chloroform, DMF or mixtures thereof. Removal of the Boc  
protecting group is accomplished by treatment of 59 with an  
acid catalyst such as TFA, or hydrogen chloride in a suitable  
35 insert solvent such as dichloromethane, ethanol or ethyl  
acetate and produces intermediate 60. Reaction of 60 with a

suitable opti nally protected guanylate reagent such as bis-Cbz-S-methyl is thiourea in an inert s lvent such as DMF or THF in the presence of a base such as triethylamine provides the protected intermediate 61. Swern oxidation of  
5 61 with a mixture of oxalyl chloride, DMSO, triethylamine in dichloromethane at about -78°C to room temperature gives the ketone intermediate 62. Finally, deprotection of 62 with anhydrous hydrogen fluoride optionally in the presence of anisole at about -78°C to room temperature produces the  
10 desired perfluoroalkylargininyl ketone 63.

Figure 12 depicts a reaction scheme for the preparation of intermediate 69 which is used in the synthesis of compounds of formula (I). Intermediate 69 may be coupled to intermediates such as 8 and 16 according to reaction schemes  
15 analogous to those depicted in Figures 4 and 5.

### 3. Selection of Preferred Compounds

The compounds of the present invention are screened for their ability to inhibit thrombin, plasmin, tissue plasminogen activator (t-PA), activated protein C (aPC), chymotrypsin, and  
20 trypsin as set forth below. Certain of the preferred compounds are distinguished by their ability to inhibit thrombin, while not substantially inhibiting plasmin, t-PA, aPC, chymotrypsin, and trypsin. With respect to thrombin and  
25 the other enzymes and as used herein, the term "not substantially inhibiting" means that the IC<sub>50</sub> (or K<sub>i</sub>) for plasmin, t-PA, aPC, chymotrypsin, and/or trypsin for a given compound is greater than or equal to its IC<sub>50</sub> (or K<sub>i</sub>, respectively) for thrombin.

30 With respect to compounds within the present invention that inhibit members within the trypsin/chymotrypsin family, including trypsin, chymotrypsin, elastase, and serine proteases involved in the coagulation cascade, "not specifically inhibiting" means the IC<sub>50</sub> or K<sub>i</sub> for the target

enzyme is less than or equal to the  $IC_{50}$  or  $K_i$  for non-target enzymes contacted with the inhibitor.

For screening compounds using these assays, the compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from 0 to 100 micromolar. In the assays for thrombin, plasmin, t-PA, aPC, chymotrypsin, and trypsin, a chromogenic synthetic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The  $IC_{50}$  of a compound of the present invention is determined from the rate of substrate turnover caused by the specific enzyme being measured.  $IC_{50}$  is the concentration of test compound which gives 50% inhibition of the rate of substrate turnover. Likewise, the  $K_i$  of a compound of the present invention is determined from the rate of substrate turnover caused by the specific enzyme being measured at various enzyme concentrations. Examples A, B and C provide an exemplars of in vitro assays used to select the compounds of the present invention.

Certain of the preferred compounds of the present invention have a  $K_i$  of about 0.001 to about 200 nM in the thrombin assay. Especially preferred compounds have a  $K_i$  of about 0.001 to about 50 nM. The more especially preferred compounds have a  $K_i$  of about 0.001 to about 10 nM.

Certain of the preferred compounds of the present invention have a  $IC_{50}$  for plasmin, t-PA, aPC, chymotrypsin, and trypsin which is at least 10 times greater than its  $IC_{50}$  for thrombin. Especially preferred compounds have an  $IC_{50}$  for plasmin, t-PA, aPC, chymotrypsin, and trypsin which is about 20 to about 100,000 times greater than its  $IC_{50}$  for thrombin. More especially preferred compounds have an  $IC_{50}$  for plasmin, t-PA, aPC, chymotrypsin, and trypsin which is about 100 to about 1,000,000 times greater than its  $IC_{50}$  for thrombin. In the event that a compound of the present



invention has an IC<sub>50</sub> with respect to plasmin, t-PA, aPC, chymotrypsin, or trypsin which is greater than the highest concentration of compound tested, the highest concentration of compound tested is considered to be the reported IC<sub>50</sub>.

5

#### 4. Pharmaceutical Compositions

In another aspect, the present invention encompasses pharmaceutical compositions prepared for storage or administration which comprise a therapeutically effective amount of a compound of the present invention in a pharmaceutically acceptable carrier.

The "therapeutically effective amount" of a compound of the present invention will depend on the route of administration, the type of mammal being treated, and the physical characteristics of the specific mammal under consideration. These factors and their relationship to determining this amount are well known to skilled practitioners in the medical arts. This amount and the method of administration can be tailored to achieve optimal efficacy but will depend on such factors as weight, diet, concurrent medication and other factors which, as noted, those skilled in the medical arts will recognize.

The "therapeutically effective amount" of the compound of the present invention can range broadly depending upon the desired affects and the therapeutic indication. Typically, dosages will be between about 0.01 mg/kg and 100 mg/kg body weight, preferably between about 0.01 and 10 mg/kg, body weight.

"Pharmaceutically acceptable carriers" for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A.R. Gennaro edit. 1985). For example, sterile saline and phosphate-buffered saline at physiological pH may be used. Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. For example, sodium benzoate,

sorbic acid and esters of p-hydroxybenzoic acid may be added as preservatives. *Id.* at 1449. In addition, antioxidants and suspending agents may be used. *Id.*

5 The pharmaceutical compositions of the present invention may be formulated and used as tablets, capsules or elixirs for oral administration; suppositories for rectal administration; sterile solutions and suspensions for injectable administration; and the like. The dose and method of administration can be tailored to achieve optimal efficacy but  
10 will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

When administration is to be parenteral, such as intravenous on a daily basis, injectable pharmaceutical  
15 compositions can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example; water, saline, dextrose, mannitol, lactose, lecithin, albumin, sodium  
20 glutamate, cysteine hydrochloride, or the like. In addition, if desired, the injectable pharmaceutical compositions may contain minor amounts of nontoxic auxiliary substances, such as wetting agents, pH buffering agents, and the like. If  
25 desired, absorption enhancing preparations (e.g., liposomes) may be utilized.

#### 5. Utility.

Certain compounds of the present invention when made and selected as disclosed are useful as potent inhibitors of  
30 thrombin *in vitro* and *in vivo*. As such, these compounds are useful as *in vitro* diagnostic reagents to prevent the clotting of blood and as *in vivo* pharmaceutical agents to prevent thrombosis in mammals suspected of having a condition characterized by abnormal thrombosis.

35 The compounds of the present invention are useful as *in vitro* diagnostic reagents for inhibiting clotting in blood

drawing tubes. The use of stoppered test tubes having a vacuum therein as a means to draw blood obtained by venipuncture into the tube is well known in the medical arts. See, e.g., Kasten, B.L., "Specimen Collection", Laboratory Test Handbook, 2nd Edition, Lexi-Comp Inc., Cleveland pp. 16-17 (Edits. Jacobs, D.S. et al. 1990). Such vacuum tubes may be free of clot-inhibiting additives, in which case, they are useful for the isolation of mammalian serum from the blood. They may alternatively contain clot-inhibiting additives (such as heparin salts, EDTA salts, citrate salts or oxalate salts), in which case, they are useful for the isolation of mammalian plasma from the blood. The compounds of the present invention are potent inhibitors of blood clotting enzymes, such as factor Xa or thrombin, and as such, can be incorporated into blood collection tubes to prevent clotting of the mammalian blood drawn into them.

The compounds of the present invention are used alone, in combination of other compounds of the present invention, or in combination with other known inhibitors of clotting, in the blood collection tubes. The amount to be added to such tubes is that amount sufficient to inhibit the formation of a clot when mammalian blood is drawn into the tube. The addition of the compounds to such tubes may be accomplished by methods well known in the art, such as by introduction of a liquid composition thereof, as a solid composition thereof, or liquid composition which is lyophilized to a solid. The compounds of the present invention are added to blood collection tubes in such amounts that, when combined with 2 to 10 mL of mammalian blood, the concentration of such compounds will be sufficient to inhibit clot formation. Typically, the required concentration will be about 1 to 10,000 nM, with 10 to 1000 nM being preferred.

Certain of the compounds of the present invention are useful as a pharmaceutical agent for preventing thrombosis in a mammal suspected of having a condition characterized by abnormal thrombosis.

Conditions characterized by abnormal thrombosis are well known in the medical arts and include those involving the arterial and venous vasculature of mammals. With respect to the coronary arterial vasculature, abnormal thrombosis (thrombus formation) characterizes the rupture of an established atherosclerotic plaque which is the major cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA). With respect to the venous vasculature, abnormal thrombosis characterizes the condition observed in patients undergoing major surgery in the lower extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombosis further characterizes disseminated intravascular coagulopathy which commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.

The present invention includes methods for preventing a condition in a mammal suspected of having a condition characterized by abnormal thrombosis, comprising administering to said mammal a therapeutically effective amount of a compound or a pharmaceutical composition of the present invention.

The compounds or pharmaceutical compositions of the present invention are administered *in vivo*, ordinarily in a mammal, preferably in a human. In employing them *in vivo*, the compounds or pharmaceutical compositions can be administered to a mammal in a variety of ways, including orally, parenterally, intravenously, subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing

a variety of dosage forms. Administration is preferably parenteral, such as intravenous on a daily basis. Alternatively, administration is preferably oral, such as by tablets, capsules or elixers taken on a daily basis.

5 In practicing the methods of the present invention, the compounds or pharmaceutical compositions of the present invention are administered alone or in combination with one another, or in combination with other therapeutic or in vivo diagnostic agents.

10 As is apparent to one skilled in the medical art, a therapeutically effective amount of the compounds or pharmaceutical compositions of the present invention will vary depending upon the age, weight and mammalian species treated, the particular compounds employed, the particular mode of  
15 administration and the desired affects and the therapeutic indication. Because these factors and their relationship to determining this amount are well known in the medical arts, the determination of therapeutically effective dosage levels, the amount necessary to achieve the desired result of  
20 preventing thrombosis, will be within the ambit of one skilled in these arts. Typically, administration of the compounds or pharmaceutical composition of the present invention is commenced at lower dosage levels, with dosage levels being increased until the desired effect of preventing in vivo  
25 thrombosis is achieved which would define a therapeutically effective amount. For the compounds of the present invention, alone or as part of a pharmaceutical composition, such doses are between about 0.01 mg/kg and 100 mg/kg body weight, preferably between about 0.01 and 10 mg/kg, body weight.

30 Certain compounds of the present invention have utility as inhibitors of proteases within the trypsin/chymotrypsin class of enzymes. Members of that class include, but are not limited to, elastase, chymotrypsin, and the serine proteases trypsin, thrombin, factor Xa, and factor VIIa. With respect  
35 to the inhibitors within the present invention directed at serine proteases acting within the coagulation cascade, e.g.

inhibitors of thrombin, factor Xa and factor VIIa, such have in vitro and in vivo utilities as provided hereinabove for thrombin inhibitors.

Elastase has been implicated in a variety of conditions, including pulmonary hypertension (Rabinovitch, M., *Acta Paediatr. Jpn* 37:657-666 (1995)), idiopathic pulmonary fibrosis, rheumatoid arthritis, adult respiratory distress syndrome, cystic fibrosis, and other inflammatory diseases and conditions (Doring, G., *Am. J. Respir. Crit. Care Med.* 150:S114-S117 (1994)). Inhibition of elastase was shown to prevent or retard progression of pulmonary hypertension (Rabinovitch). Thus, inhibitors of the present invention directed toward elastase are useful as pharmaceutical compositions for the inhibition of elastase in conditions where elastase activity is associated with a pathological condition.

Elevated levels of chymotrypsin and trypsin are associated with the pathological effects resulting from pancreatitis (see US 5,534,498). Animal studies of chemically-induced pancreatitis suggest that the disorder is rooted in the inability of pancreatic acinar cells to excrete digestive proenzymes, resulting in activation of trypsinogen to trypsin by lysosomal hydrolases within the cell. The amount of trypsin produced exceeds protective levels of protease inhibitor normally available.

The elevated levels of trypsin then cause activation of the other digestive enzymes co-localized with trypsin in the lysosome, such as chymotrypsin. The net effect of the enzyme activation is pancreatitis, which is characterized by damage to the pancreas and surrounding tissues from autodigestion of the cells by the various digestive enzymes. These activated digestive enzymes also cause edema, interstitial hemorrhage, vascular damage, coagulation necrosis, fat necrosis and parenchymal cell necrosis.

Inhibitors of the present invention directed toward either trypsin or chymotrypsin, or other members of the

trypsin/chymotrypsin family that contribute to the deleterious effects of pancreatitis, are useful for the prevention and treatment of pancreatitis in mammals.

In addition to the *in vivo* utilities, inhibitors of the present invention also find utility *in vitro*. Inhibitors of enzymes within the coagulation cascade are useful inhibitors of blood coagulation *in vitro*, as described hereinabove. Inhibitors of other enzymes within the trypsin/chymotrypsin family, including trypsin, chymotrypsin, and elastase, are useful reagents in *in vitro* assays designed to measure the activity of such enzymes.

For instance, to determine or confirm the presence of active trypsin, chymotrypsin, or elastase in a sample, the activity of the enzyme in the sample is determined in the presence and absence of the specific inhibitor (which may be labeled using a radioactive or other detectable label). Lower activity measured in the presence of inhibitor as compared to in the absence of inhibitor demonstrates inhibition of the enzyme and, thus, its presence in the sample.

Similarly, the level of activity of an enzyme present in a sample is determined by adding inhibitor to the sample in a range of titrating doses, and calculating activity of the enzyme at each escalating dose of inhibitor. The concentration of inhibitor that completely inhibits the enzyme in the assay, along with knowledge of the assay parameters and characteristic of enzyme inhibition, allows one to calculate the activity of the enzyme in the sample.

The level of chymotrypsin measured in stool samples *in vitro* is used as an indicator of pancreatitis (Riedel, L. et al. Gut 32:321-324 (1991); Chari, S., Trop. Gastroenterol. 11:144-147 (1990)). Chymotrypsin inhibitors of the present invention are useful in such assays to evaluate the level of active chymotrypsin in such a sample, according to protocols such as those outlined hereinabove.

An additional use of the inhibitors of the present invention is their use to quench enzymatic reactions effected

by the target enzyme. Thus, to control or prevent digestion of a sample with trypsin or chymotrypsin, an inhibitor of trypsin or chymotrypsin, respectively, is added to inhibit the target enzyme and, thus, control or prevent digestion by that enzyme.

Certain compounds of the present invention can also be useful inhibitors of elastase, and are therefore useful pharmaceutical agents for the control of inflammation.

To assist in understanding, the present invention will now be further illustrated by the following examples. These examples as they relate to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

Examples 1 to 11 describe preparation of compound 11 of Figure 1.

Examples 12 to 14 describe preparation of compound 19 of Figure 2.

Examples 15 to 19 describe preparation of intermediate 25 of Figure 3.

Examples 20 to 22 describe preparation of compound 28 of Figure 4.

Examples 24 to 30 describe preparation of compound 41 of Figure 6.

Examples 31 to 33 describe preparation of compound 44 of Figure 7.



EXAMPLESExample 1Preparation of N-(2-Hydroxy-4-methylphenyl)phthalimide (1)

A solution of phthalic anhydride (14.81 g, 0.10 mole) and 6-amino-m-cresol (12.32 g, 0.10 mole) in 200 mL of acetic acid was stirred at ambient temperature for 30 minutes and then refluxed for 6 hours. The solution was cooled and poured into 500 mL ice-water with rapid stirring and the resultant solid was collected by suction filtration, washed with ice-cold water several times and air dried overnight to afford 24.1 g (95% yield) of product 1 as a light brown solid, mp 251-254°C. TLC (silica ; EtOAc), R<sub>f</sub> = 0.68.

Example 2Preparation of N-(2-Allyloxy-4-methylphenyl)phthalimide (2)

To a mixture of 1 (23.78 g, 0.0939 mole; see Example 1) and anhydrous K<sub>2</sub>CO<sub>3</sub> (15.62 g, 0.113 mole) in 190 mL of anhydrous acetonitrile was added allyl bromide (17.04 g, 0.141 mole, 12.2 mL). The mixture was stirred under N<sub>2</sub> at room temperature for 15 minutes and then refluxed for 5 hours. The mixture was cooled, the solids were filtered and then washed with fresh acetonitrile/ethyl acetate, and the filtrate was evaporated. The residue was dissolved in 500 mL of ethyl acetate, extracted with 50 mL portions of saturated NaHCO<sub>3</sub> solution (2x), water, brine and dried over anhydrous MgSO<sub>4</sub>. Filtration and removal of solvent afforded 26.76 g (97% yield) of product 2 as a light brown solid, mp 122-124°, judged pure by TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>), R<sub>f</sub> = 0.45.

Example 3Preparation of N-(2-Hydroxy-3-allyl-4-methylphenyl)phthalimide (3)

A solution of 2 (2.93 g, 0.010 mole; see Example 2) in 6 mL of N,N-dimethylaniline under N<sub>2</sub> was heated in an oil bath to 185°C (pot temperature) for 1.5 hours and then cooled. The reaction mixture was poured onto a flash silica gel column

and eluted directly with a hexane, CH<sub>2</sub>Cl<sub>2</sub>:2,1 to CH<sub>2</sub>Cl<sub>2</sub> gradient solvent system to afford 2.79 g (95% yield) of product 3 as a colorless solid, mp 178-180°C. TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>), R<sub>f</sub> = 0.25. Scale-up to 23.7 g and a reaction time of 5 2.5 hours afforded the product in 91% yield, identical in all respects with the above material.

#### Example 4

#### Preparation of N-(2-Benzoyloxy-3-allyl-4-methylphenyl)phthalimide (4)

To a mixture of 3 (6.60 g, 0.0225 mole; see Example 3) and anhydrous K<sub>2</sub>CO<sub>3</sub> (3.73 g, 0.027 mole) in 45 mL of anhydrous acetonitrile was added benzyl bromide (5.00 g, 0.0293 mole, 3.49 mL). The mixture was stirred under N<sub>2</sub> at 15 room temperature for 15 minutes and then refluxed for 19 hours. The mixture was cooled and filtered; the collected solids were washed with fresh acetonitrile and the filtrate was evaporated. The residue was dissolved in 300 mL of ethyl acetate and extracted with 50 mL portions of saturated 20 NaHCO<sub>3</sub> (2x), water, brine and dried over anhydrous MgSO<sub>4</sub>. Filtration, solvent removal and recrystallization from 1-chlorobutane afforded product as a solid. The filtrate residue was flash chromatographed on silica gel, eluting with a hexane, CH<sub>2</sub>Cl<sub>2</sub>:2,1 to CH<sub>2</sub>Cl<sub>2</sub> gradient to afford a total of 25 6.99 g (81% yield) 4 as a colorless solid, mp 107-109°C. TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>, Hexane: 2,1), R<sub>f</sub> = 0.30.

Example 5Preparation of 2-Benzylloxy-3-allyl-4-methylaniline (5)

To a solution of 4 (7.67 g, 0.020 mole; see Example 4) in 100 mL of ethanol was added hydrazine monohydrate (1.22 g, 0.024 mole, 1.18 mL). The reaction was stirred at room temperature under N<sub>2</sub> for 15 minutes, refluxed for 27 hours, cooled and filtered, and then the filtrate was evaporated. The residue was purified by flash chromatography on silica gel, using a hexane, ethyl acetate: 9,1 to 4,1 gradient and afforded 5.03 g (99% yield) of product 5 as a viscous light red-brown oil. TLC (silica; hexane, ethyl acetate: 2,1), R<sub>f</sub> = 0.45.

Example 6Preparation of N-Benzylsulfonanilide, 2-benzylloxy-3-allyl-4-methyl- (6)

To a solution of 5 (7.54 g, 0.0298 mole; see Example 5) and 4-methylmorpholine (9.04 g, 0.0894 mole, 9.83 mL) in 300 mL of anhydrous acetonitrile at 0°C under N<sub>2</sub> was added  $\alpha$ -toluene-sulfonyl chloride (11.36 g, 0.0596 mole) in one portion. The solution was stirred at 0°C for 1 hour and allowed to warm to and stir at ambient temperature for 19 hours. The solution was filtered and evaporated. The crude residue was purified by flash chromatography on silica gel using a hexane, CH<sub>2</sub>Cl<sub>2</sub>: 4,1 to CH<sub>2</sub>Cl<sub>2</sub> gradient system and afforded 12.05 g (99% yield) of product 6 as waxy reddish solid, mp 62-63°C. TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>), R<sub>f</sub> = 0.45.

Example 7Preparation of 2-Benzylloxy-3-(N-benzylsulfonylamino)-6-methylphenylacetaldehyde (7)

A stream of ozone gas was introduced into a solution of 6 (10.71 g, 0.0263 mole; see Example 6) dissolved in 260 mL of methanol at -78°C until a pale blue color persisted. After 3 minutes, a stream of dry N<sub>2</sub> was bubbled through the solution to discharge excess ozone. Dimethyl sulfide (3.27

g, 0.0526 mole, 3.86 mL) was added, the solution was stirred at -78°C for 5 minutes, and was allowed to warm to and then stir at ambient temperature for 14 hours. The solvent was evaporated to afford a residue which was dissolved in 350 mL of diethyl ether, extracted with 50 mL portions of water (2x), brine, dried over anhydrous MgSO<sub>4</sub>, filtered and evaporated to afford 11.73 g of a viscous oil. This oil was dissolved in 245 mL acetone and 75 mL of 1N HCl was added. The solution was stirred at room temperature for 14 hours, refluxed for 30 minutes, cooled and evaporated. The residue was dissolved in 600 mL of diethyl ether and extracted with 50 mL portions of saturated NaHCO<sub>3</sub>, water (2x), brine, and dried over MgSO<sub>4</sub>. Filtration and solvent removal afforded 9.71 g (98% yield) of product 7 as a viscous yellow oil. TLC (silica; hexane, ethyl acetate: 2,1), R<sub>f</sub> = 0.25.

#### Example 8

#### Preparation of 2-Benzoyloxy-3-(N-benzylsulfonylamino)-6-methylphenylacetic acid (8)

To a solution of 7 (9.70 g, 0.0237 mole; see Example 7) in 250 mL of acetone at 0°C was added Jones' reagent rapidly dropwise over a 30 minute period until a brown color persisted. The ice bath was removed and the mixture was allowed warm to room temperature. 2-Propanol (15 mL) was added and the mixture was stirred for 10 minutes. The mixture was diluted with 500 mL of ethyl acetate and filtered with suction through a pad composed of a homogeneous 1:1 mixture of celite and flash-grade silica gel. Elution with an additional 1 liter portion of acetone, ethyl acetate 1,1, followed by solvent removal afforded 10.30 g (-Quantitative crude yield) of product 8 as a sticky yellow foam, judged nearly pure by TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>, EtOH: 9,1), R<sub>f</sub>=0.53. A portion recrystallized from diethyl ether and hexane afforded a colorless solid, mp 132-134°C.

Example 9Preparation of N-[2-Benzylloxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-N<sup>9</sup>-nitro-L-argininal Ethyl Cyclol (9)

To a solution of 8 (9.16 g, 0.02153 mole; see Example 8) in 215 mL of anhydrous acetonitrile at room temperature under N<sub>2</sub> was added N<sup>9</sup>-nitro-L-argininal ethyl cyclol (6.92 g, 0.0258 mole), 1-hydroxybenzotriazole hydrate (3.30 g, 0.0215 mole) and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (6.19 g, 0.0323 mole). After stirring the mixture for 15 minutes, N,N-diisopropylethylamine (11.13 g, 0.0861 mole, 15.0 mL) was added and the resultant clear light brown solution was stirred at room temperature for 22 hours. The solvent was evaporated, the residue was slowly dissolved in 700 mL of ethyl acetate and was extracted with 50 mL portions of 1N HCl (3x), saturated NaHCO<sub>3</sub> (2x), water (2x), brine (2x), dried over MgSO<sub>4</sub>, filtered and evaporated. The crude product was purified by flash chromatography on silica gel using a CH<sub>2</sub>Cl<sub>2</sub>, EtOH: 99,1 to 95,5 gradient system to afford 9.49 g (69% yield) of product 9 as a light yellow foam. TLC (silica; ethyl acetate), R<sub>f</sub> = 0.50.

Example 10Preparation of N-[2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-L-argininal Ethyl Cyclol Acetate Salt (10)

To a solution of 9 (9.27 g, 0.0145 mole; see Example 9) in 370 mL of ethanol, acetic acid, water: 4,1,1 was added 10% Pd/C (9.3 g). The mixture was hydrogenated at 60 PSI on a Parr shaker apparatus for 20 hours. The mixture was filtered and the solvents were evaporated. The residue was dissolved in ethanol, acetonitrile 1,1, reevaporated, and pumped at < 1 mm Hg on a vacuum pump for 24 hours to afford 7.23 g (89% yield) of product 10 as a yellow foam. TLC (silica; CH<sub>2</sub>Cl<sub>2</sub>, methanol, conc NH<sub>4</sub>OH: 20, 10, 2), R<sub>f</sub> = 0.50.

Example 11Preparation of N-(2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl)-L-argininal Trifluoroacetate Salt (11)

Compound 10 (1.01 g, 1.78 mmole; see Example 10) was  
5 added to a solution of 35 mL of 9.4 M HClO<sub>4</sub> at 0°C. The  
starting material slowly dissolved over 30 minutes and the  
reaction mixture was then frozen by immersion in a dry-ice  
acetone slurry. After warming to -15°C, the reaction mixture  
was adjusted to pH 5 by using a 10 M NaOH solution followed  
10 by a saturated NaOAc solution. During the neutralization  
process, the temperature was maintained at <10°C. The  
solution was stored at 4°C for one hour, then the solids were  
filtered. The filtrate was purified by reverse phase HPLC on  
a 50 X 300 mm C18 column using a 0 to 28% gradient of  
15 acetonitrile/water (containing 0.1% trifluoroacetic acid)  
over one hour to afford 0.66 g (63% yield) of product 11 as a  
colorless, amorphous solid. RP/HPLC analysis showed three  
peaks for the product. Mass spectrometry confirmed the  
theoretical molecular weight of 475.

20

Example 12Preparation of N-(2-Methoxy-3-allyl-4-methylphenyl)phthalimide (12)

To a mixture of 3 (6.00 g, 0.0205 mole; see Example 3)  
25 and anhydrous K<sub>2</sub>CO<sub>3</sub> (3.39 g, 0.025 mole) in 41 mL of  
anhydrous acetonitrile was added iodomethane (5.81 g, 0.0409  
mole, 2.54 mL). The mixture was stirred under N<sub>2</sub> at room  
temperature for 15 minutes and then refluxed for 17 hours.  
The mixture was cooled and filtered, the collected solids  
30 were washed with fresh acetonitrile and the filtrate was  
evaporated to afford 5.08 g (81% yield) of pure 12 as a  
colorless solid. TLC (silica; Hexane, ethyl acetate:1,1), R<sub>f</sub>  
= 0.50.


Example 13Preparation of N-[2-Methoxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-L-argininal Ethyl Cyclohexyl Acetate Salt (18)

5       Compound 18 was prepared from compound 12 by following an analogous six-step protocol outlined above for the preparation of compound 10 from compound 4 (hydrazinolysis, benzylsulfonation, ozonolysis, oxidation, coupling and hydrogenation; see Examples 5 to 10).

10

Example 14Preparation of N-[2-Methoxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-L-argininal Trifluoroacetate Salt (19)

15       Compound 18 (1.00 g, 1.73 mmole) was added to a solution of 72 mL of 3N HCl and 5 mL of acetonitrile. The mixture was stirred at room temperature for 2 hours, 10.0 mL of 6N HCl was added, and reaction was continued for a further 3.5 hours. The solution was filtered and purified by reverse phase HPLC on a 50 X 300 mm C18 column using a 5 to 30%  
20       gradient of acetonitrile/water (containing 0.1% trifluoroacetic acid) over one hour and afforded 0.77 g (74% yield) of product 19 as a colorless, amorphous solid. RP/HPLC analysis showed three peaks for the product. Mass spectrometry confirmed the theoretical molecular weight of  
25       489.


Example 15Preparation of N-(t-butoxycarbonyl)-3-(3-pyridyl)-L-alanine methyl ester (21)

To a solution of N-(t-butoxycarbonyl)-3-(3-

pyridyl)alanine (20) (5.0 g, 18.8 mmole) in methanol (100 mL) was added thionyl chloride (2M solution in dichloromethane, 66 mL, 132 mmole). The resulting solution was stirred overnight at ambient temperature. The methanol was removed under reduced pressure to a minimum volume and ethyl acetate (100 mL) was added. The resulting white precipitate was collected in a fritted funnel. To a solution of the collected precipitate in a mixture of tetrahydrofuran/water (40 mL each) was added di-tert-butyl dicarbonate (4.8 g, 21.99 mmole) and sodium carbonate (1.95 g, 18.4 mmole). After stirring for 12 hours at ambient temperature, the reaction mixture was diluted with ethyl acetate (40 mL) and washed with a solution of saturated sodium bicarbonate (25 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum to give crude product. This product was subjected to flash column chromatography on silica gel (230-400 mesh) using a 8x52 cm column and eluting with a 10:90 mixture of ethyl acetate/hexane followed by a 60:40 mixture of ethyl acetate/hexane. 4 g (74%) of the title compound was obtained as an oil. Thin-layer chromatography (silica gel; ethyl acetate),  $R_f$  = 0.68.

#### Example 16

Preparation of N-(t-butoxycarbonyl)-3-([(R,S)-3-piperidyl])-L-alanine methyl ester, acetate salt (22)




A solution of the compound of Example 15 (5 g, 17.8 mmole) in ethanol (24 mL), acetic acid (6 mL) and water (6 mL) was hydrogenated over platinum oxide (500 mg) at 45 psi for three hours. The catalyst was filtered off and the



filtrate concentrated under vacuum to an oily residue (6.89 g) which was used in the next step (Example 17) without further purification. Thin-layer chromatography yielded two spots corresponding to two diastereomers with  $R_f$  values of 0.16 and 0.26, respectively (silica gel; 4:1:1 n-butanol/acetic acid/water).

#### Example 17

Preparation of N-(t-butoxycarbonyl)-3-[(R,S)-3-piperidyl-(N-guanidino (bis-benzyloxycarbonyl))]-L-alanine methyl ester (23)




To a solution of the compound of Example 16 (6.89 g, 19.9 mmole) in tetrahydrofuran (80 mL) was added S-methylisothiourea bis-benzyloxycarbonyl (7.13 g, 19.9 mmole) followed by N-methylmorpholine (4.37 mL), and the reaction mixture was stirred at ambient temperature for 18 hours. The reaction mixture then was concentrated under vacuum and the resulting residue was dissolved in ethyl acetate (100 mL) and washed with 1N sodium bisulfate and saturated sodium chloride (50 mL each). After drying over anhydrous sodium sulfate, the solvents were removed under vacuum; the crude title compound was subjected to flash column chromatography on silica gel (230-400 mesh) using a 8x52 cm column and eluting with 1:9 ethyl acetate/hexanes (two column volumes) followed by 1:1 ethyl acetate/hexanes. 2.75 g the title compound was obtained as a mixture of two diastereomers. Thin-layer chromatography gave two spots with  $R_f$  values of 0.57 and

0.62, respectively (silica gel; 1:1 ethyl acetate/ hexanes).

Example 18


Preparation of N-(t-butoxycarbonyl)-3-[(R,S)-3-piperidyl-(N-guanidino (bis-benzoyloxycarbonyl))]-L-alaninol (24)



- To a stirred solution of the compound of Example 17 (2.23 g, 3.7 mmole) in absolute ethanol (8 mL) and anhydrous tetrahydrofuran (4 mL) was added calcium chloride (844 mg, 7.6 mmole) and sodium borohydride (575 mg, 15.2 mmole). After stirring 12 hours at ambient temperature, the reaction mixture was concentrated under vacuum and the resulting residue was partitioned between ethyl acetate and 1N sodium bisulfate (10 mL each). The two layers were separated; organic layer was washed twice more with 1N sodium bisulfate, dried over anhydrous sodium sulfate and concentrated under vacuum gave a residue. Flash column chromatography of the residue on silica gel (230-400 mesh) using a 5.5x45 cm column and eluting with ethyl acetate gave 1.3 g of the title compound as a white foam. Thin layer chromatography yielded two spots corresponding to two diastereomers with  $R_f$  values of 0.18 and 0.27, respectively (silica gel; 1:1 ethyl acetate/ hexanes).

Example 19

Preparation of 3-[(R,S)-3-piperidyl-(N-guanidino(bis-benzoyloxycarbonyl))]-L-alaninol, hydrochloride salt (25)



5

The compound of Example 18 (290 mg, 0.57 mmole) was treated with 2.5 N anhydrous hydrochloric acid in ethyl acetate (2.0 mL) at ambient temperature for one hour. The solvent was removed under vacuum to a sticky-white solid (260 mg). This solid was used in the next step (Example 9) without further purification. <sup>1</sup>H NMR spectrum taken in CD<sub>3</sub>OD showed no t-butoxycarbonyl protons at 1.4 ppm.

Example 20

Preparation of N-[2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenyl-acetyl]-3-[(R,S)-3-piperidyl-(N-guanidino(bis-benzoyloxycarbonyl))]l-L-alaninol (26)

5 To a suspension of the Compound of Example 19 (25) (2.06 g, 4.08 mmole) in acetonitrile (22 mL) is added successively the compound of Example 8 (8) (2.36 g, 5.56 mmole), EDC (1.12 g, 5.84 mmole), 1-hydroxybenzotriazole hydrate (979 mg, 6.39 mmole), and N-methylmorpholine (3 mL, 10 27.80 mmole). The solution is stirred at ambient temperature for about twenty hours. The solvent is removed under vacuum and the resulting residue is dissolved in a 9:1 mixture of dichloromethane/isopropanol (40 mL) and washed two times each with 15 mL portions of 1N sodium bisulfate, saturated sodium 15 bicarbonate and saturated sodium chloride. The organic layer is dried over anhydrous sodium sulfate and concentrated under vacuum to give the title compound. The product is a mix of diastereomers at P1 and has a calculated molecular weight of 875.05.

20

Example 21

Preparation of N-[2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenyl-acetyl]-3-[(R,S)-3-piperidinyl-(N-guanidino)]l-L-alaninol (27)


25 The compound of Example 20 (26) (1.97 g, 2.25 mmole) is subjected to catalytic hydrogenation in methanol (100 mL) and acetic acid (10 mL in the presence of 10% palladium on carbon (185 mg) at 30 psi for about 2.5 hours. The catalyst is filtered. The filtrate after concentration under vacuum 30 yields the title compound as two diastereomers with a calculated molecular weight of 577.7.

Example 22Preparation of N-(2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenyl-acetyl)-3-[3-piperidinyl-(N-guanidino)-L-alaninal and separation of diastereomers (28a and 28b)

- 5 To a chilled solution of the compound of Example 21 (27) (0.81 g, 1.4 mmole) in dimethylsulfoxide and toluene (15 mL each) is added dichloroacetic acid (567  $\mu$ L, 6.9 mmole), followed by EDC (2.68 g, 14 mmole) at one minute later. The reaction mixture is stirred for about 5 minutes at 0°C, then  
10 about 85 minutes at ambient temperature, and then is quenched with 60 mL water. The water layer is extracted twice with diethyl ether (10 mL portions) and subjected to HPLC using a 47 X 300 mm reverse phase column containing a C-18 resin  
15 comprised of 10 micron-size gel particles with a 300 angstrom pore size. The column is eluted with a gradient ranging from 15% to 30% acetonitrile in water (containing 0.1% trifluoroacetic acid). The HPLC fractions will yield fast moving and slow moving peaks containing the two diastereomers of the title compound. The fractions containing each  
20 diastereomer when pooled, then lyophilized, will give the two diastereomers of the title compound, (28a and 28b).

Example 23Preparation semicarbazid-4-yl diphenylmethane.

- 25 trifluoroacetate salt


Step 1:

- 30 A solution of carbonyldiimidazole (16.2 g, 0.10 mole) in 225 mL of dimethylformamide was prepared at room temperature and allowed to stir under nitrogen. A solution

of t-butyl carbazate (13.2 g, 0.100 moles) in 225 mL dimethylformamide was then added dropwise over a 30 minute period. Next, diphenylmethanamine (18.3 g, 0.10 moles) was added over a 30 minute period. The reaction was allowed to stir at room temperature under nitrogen for one hour. Water (10 mL) was added and this mixture was concentrated to about 150 mL under vacuum. This solution was poured into 500 mL water and extracted with 400 mL of ethyl acetate. The ethyl acetate phase was extracted two times each with 75 mL 1N HCl, water, saturated sodium bicarbonate and brine, and then was dried with anhydrous magnesium sulfate. The mixture was filtered and the solution was concentrated to give 29.5 g (85% yield) of 1-t-butoxycarbonyl-semicarbazid-4-yl diphenylmethane as a white foam. This material may be purified by recrystallization from ethyl acetate/hexane, but was pure enough to use directly in step 2: mp 142-143°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) delta 1.45 (s, 9H), 6.10 (dd, 2H), 6.42 (s, 1H), 6.67 (bs, 1H), 7.21-7.31 (m, 10H). Analysis calculated for C<sub>19</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>: C, 66.84; H, 6.79; N, 12.31. Found: C, 66.46; H, 6.75; N, 12.90.

Step 2:

A solution of 3.43 g (10 mmole) of 1-t-butoxycarbonyl-semicarbazid-4-yl diphenylmethane in 12.5 mL of dichloromethane was treated with 12.5 mL of trifluoroacetic acid at 0°C. The reaction mixture was allowed to stir for 30 minutes at this temperature. The reaction mixture was then added dropwise to 75 mL of diethyl ether to give a precipitate. The resulting precipitate was filtered off and washed with diethyl ether to give 2.7 g (80% yield) of the title compound; mp 182-184°C.

Example 24Preparation of 3-thioamidobenzyl-N-acetylamino malonic acid diethyl ester (35)


To a stirred solution of alpha-bromo-meta-tolunitrile (32) (45.0 g, 0.24 mole), diethyl acetamidomalonate (33) (48.0 g, 0.22 mole) and potassium iodide (3.0 g, 0.018 mole) 10 in dioxane (500 mL) was added 2.5M sodium ethoxide in ethanol (100 mL) dropwise under an argon atmosphere. After the addition was complete, the solution was refluxed for 6 hours. The reaction mixture was allowed to stand overnight at room temperature, then diluted with brine (250 mL) and water (250 15 mL), and extracted with ethyl acetate four times (1.0 L total). The combined extracts were washed with water (100 mL), 10% citric acid (100 mL), water (100 mL) and brine (2x50 mL), then dried over anhydrous magnesium sulfate and filtered; the solvent was removed under vacuum. The crude 20 residue was recrystallized from ethyl acetate and diethyl ether in two crops to yield 43.51 g (60%) of the 3-cyanobenzyl-N-acetylamino malonic acid diethyl ester (34) as yellow crystals.

H<sub>2</sub>S(g) was bubbled into a rapidly stirring solution of 25 3-cyanobenzyl-N-acetylamino malonic acid diethyl ester (44.3 g, 0.13 mmole) in pyridine (300 mL) and triethylamine (100 mL) for 40 minutes. The reaction mixture was stirred at room temperature for 16 hours, then poured into 3.0 L of water. A yellow precipitate formed immediately. The solution was 30 allowed to stand at 4°C for 4 hours, then was filtered. The crude title compound was recrystallized from ethyl acetate

and hexanes to yield 48.1 g (98%) of the title compound as yellow crystals, m.p. 183-186°C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.31 (t,  $J=7.1$  Hz, 6H), 2.06 (s, 3H), 3.70 (s, 2H), 4.29 (q,  $J=7.1$  Hz, 4H), 4.80-4.87 (m, 1H), 6.60 (s, 1H), 7.10-7.20 (m, 1H), 7.27-7.35 (m, 2H), 7.60-7.70 (m, 2H). Analysis calculated for  $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_5\text{S}$ : C, 55.72; H, 6.05; N, 7.64. Found: C, 55.55; H, 5.96; N, 7.76.

#### Example 25

#### 10 Preparation of 3-amidino-D,L-phenylalanine, dihydrochloride salt (36)



15 The compound of Example 24 (35) (48.1 g, 0.13 mmole) was dissolved in acetone (800 mL). Iodomethane (18.3 mL, 0.19 mole, 1.5 equivalents) was added, and the solution was refluxed for 30 minutes. The solution was cooled to room temperature, and the intermediate thioimide was filtered, dried and dissolved in methanol (500 mL). Ammonium acetate (14.8 g, 0.19 mole, 1.5 equivalents) was added. The reaction mixture was refluxed for 1 hour, then cooled to room temperature, and poured into ether (1.2 L). The solution was allowed to stand at 4°C for 72 hours. The crude 3-  
25 amidinobenzyl-N-acetylamino malonic acid diethyl ester was filtered, washed with ether, air dried, and then refluxed in concentrated HCl (250 mL) for 3 hours. The reaction mixture was concentrated under vacuum, diluted with water (0.5 L), and concentrated under vacuum again. These steps were  
30 repeated. The crude title compound was purified by cation-exchange (Sephadex SP-C25) using a gradient of 0-1.0N HCl as eluent to yield 10.8g (30%) of the title compound as an off-




white solid.  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ ):  $\delta$  3.14-3.29 (2H, m), 4.17 (dd,  $J=7.4, 6.2$  Hz, 1H), 7.42-7.69 (4H, m). Analysis calculated for  $\text{C}_{10}\text{H}_{13}\text{N}_3\text{O}_2 \cdot 2\text{HCl} \cdot 1.9\text{H}_2\text{O}$ : C, 38.20; H, 6.03; N, 13.36. Found: C, 38.51; H, 5.64; N, 12.89.

5

Example 26

Preparation of N-alpha-Boc-N-omega-4-methoxy-2,3,6-trimethylbenzenesulfonyl-3-amidino-D,L-phenylalanine (37)




3-amidino-D,L-phenylalanine (the compound of Example 25, (36)) (4.00 g, 13 mmole) was dissolved in 50% aqueous dioxane (20 mL). Sodium bicarbonate (3.38 g, 40 mmole) was added, followed by di-*t*-butyl dicarbonate (2.93 g, 13 mmole) in dioxane (4 mL). The reaction mixture was stirred for 18 hours at room temperature. The solution was cooled in an ice bath, and 4.0 N sodium hydroxide was added dropwise until the solution was pH 12. 4-methoxy-2,3,6-trimethylbenzenesulfonyl chloride (8.01 g, 32 mmole) in dioxane (10 mL) was added dropwise. 4.0 N sodium hydroxide was added as needed to keep the pH at 12. The ice bath was removed. After 1 hour, 1.0 N HCl was added to bring the solution to pH 7-8. The solution was diluted with an additional 50 mL of water and then was washed with ethyl acetate two times (20 mL each). The aqueous layer was acidified to pH 1.0 with 1.0 N HCl and extracted with ethyl acetate three times (100 mL total). The combined organic layers were washed with water (20 mL) and brine twice (10 mL each). The organic layer was dried over anhydrous magnesium sulfate and the solvent was removed under vacuum. The residue was dissolved in a minimum amount of

dichloromethane, then added dropwise to ether (25 mL). Solid impurities were removed by filtering and the solvent removed from the filtrate under vacuum to give 4.90 g (68% crude yield) of the title compound as an off-white foam. A 30 mg sample of the title compound was further purified by preparative thin-layer chromatograph developing with 1% acetic acid/5% isopropanol/ dichloromethane to give 9 mg of the title compound in a purer form.  $R_f = 0.16$  (1% acetic acid/5% isopropanol/dichloromethane).  $^1H$  NMR ( $CD_3OD$ ):  $\delta$  1.32 (s, 9H), 2.14 (s, 3H), 2.63 (s, 3H), 2.71 (s, 3H), 2.93 (dd,  $J=13.7, 9.3$  Hz, 1H), 3.22 (dd,  $J=13.7, 4.3$  Hz, 1H), 3.85 (s, 3H), 4.34-4.37 (m, 1H), 6.72 (s, 1H), 7.35-7.47 (2H, m), 7.69-7.75 (m, 2H).

15 Example 27

Preparation of N-alpha-Boc-N-(4-methoxy-2,3,6-trimethylbenzenesulfonyl)-3-amidino-D,L-phenylalanine-N-methyl-O-methyl-carboxamide (38)



To a stirred solution of compound of Example 26 (37) (1.00 g, 1.92 mmole), O,N-dimethyl hydroxylamine hydrochloride (375 mg, 3.85 mmole), hydroxybenzotriazole hydrate (294 mg, 1.92 mmole) and 4-methylmorpholine (1.06 mL, 9.62 mmole) in tetrahydrofuran (4 mL), cooled in an ice bath, was added EDC (406 mg, 2.12 mmole). The ice bath was removed, and the reaction mixture was stirred for 2 hours at room temperature. The reaction mixture was diluted with ethyl acetate (75 mL), washed with water, 10% citric acid, water, saturated sodium bicarbonate, and brine. The organic


25

30

layer was dried over anhydrous magnesium sulfate and the solvent was removed under vacuum. 750 mg (69%) of the title compound was isolated.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.33 (s, 9H), 2.14 (s, 3H), 2.66 (s, 3H), 2.75 (s, 3H), 2.80-2.88 (m, 1H), 3.06-3.20 (m, 4H), 3.70 (s, 3H), 3.84 (s, 3H), 4.98-5.06 (m, 1H), 5.21 (d,  $J=8.7$  Hz, 1H), 6.48 (bs, 1H), 6.58 (s, 1H), 7.30-7.34 (m, 2H) 7.60-7.68 (m, 2H), 8.11 (bs, 1H). Analysis calculated for  $\text{C}_{27}\text{H}_{38}\text{N}_4\text{O}_7\text{S} \cdot 0.5\text{H}_2\text{O}$ : C, 56.73; H, 6.88; N, 9.80. Found: C, 56.97; H, 6.66; N, 9.43.

### Example 28

Preparation of N-alpha-Boc-N-O-4-methoxy-2,3,6-trimethylbenzenesulfonyl-D,L-3-amidinophenylalaninal (39)




To a stirred solution of  $\text{LiAlH}_4$  (2.00 mL of a 1.0 M solution in tetrahydrofuran, 1.24 mmole) in tetrahydrofuran (8 mL), cooled in a dry ice/acetone bath, the compound of Example 27 (38) (0.75 g, 1.9 mmole in tetrahydrofuran (5 mL)) was added dropwise. The cooling bath was removed and the reaction mixture was allowed to warm to  $5^\circ\text{C}$ . The reaction mixture was re-cooled in the dry ice acetone bath and quenched with 3.0 mL of a 1:2.7 wt./wt. solution of potassium bisulfate in water. The reaction mixture was allowed to warm to room temperature, stirred for 3 hours, filtered and concentrated under vacuum. The residue was dissolved in ethyl acetate (20 mL), and washed with 10% citric acid (2 mL), water (2 mL), saturated sodium bicarbonate (2 mL) and brine (2 mL). The organic layer was dried over anhydrous magnesium sulfate and the solvent was removed under vacuum to

yield 580 mg (86%) of the title compound.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  
 $\delta$  1.31 (s, 9H), 2.07 (s, 3H), 2.57 (s, 3H), 2.67 (s,  
3H), 2.90-3.17 (2H, m), 3.77 (s, 3H), 4.33-4.40 (1H, m), 5.02-  
5.08 (1H, m), 6.48 (1H, s), 7.23-7.31 (2H, m), 7.50-7.62 (2H,  
5 m), 7.94, (1H, bs), 8.05 (1H, bs), 9.55 (1H, s). Analysis  
calculated for  $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_6\text{S}\cdot 0.5\text{H}_2\text{O}$ : C, 58.58; H, 6.69;  
N, 8.20. Found: C, 58.57; H, 6.72; N, 7.98.

#### Example 29

10 Preparation of N-alpha-Boc-N-O-4-methoxy-2,3,6-  
trimethylbenzenesulfonyl-D,L-3-amidinophenylalaninal-  
semicarbazonyl-4-N-diphenylmethane (40)




15

The compound of Example 28 (39) (0.58 g, 1.9 mmole), the  
compound of Example 23 (410 mg, 1.15 mmole) and sodium  
acetate trihydrate (188 mg, 1.38 mmole) were refluxed in 75%  
aqueous ethanol (10 mL) for 1 hour. After the reaction  
20 mixture was cooled to room temperature, it was diluted with  
ethyl acetate (50 mL), washed with 1.0N HCl (5 mL), water (5  
mL), saturated sodium bicarbonate (5 mL) and brine (2x5 mL),  
and dried over anhydrous magnesium sulfate. The solvent was  
removed under vacuum to yield 750 mg (89% yield) of the title  
25 compound as an off-white foam. Analysis calculated for  
 $\text{C}_{39}\text{H}_{46}\text{N}_6\text{O}_6\text{S}\cdot 1.0\text{H}_2\text{O}$ : C, 62.88; H, 6.49; N, 11.28. Found:  
C, 63.14; H, 6.35 N, 11.10. Calculated molecular weight  
was 726.90.

Example 30

Preparation of N- $\omega$ -4-methoxy-2,3,6-trimethylbenzene sulfonyl-D,L-3-amidinophenylalaninal-semicarbazonyl-4-N-diphenylmethane, trifluoroacetate salt (41)



The compound of Example 29 (40) (750 mg, 1.9 mmole) was treated with 50% trifluoroacetic acid/dichloromethane (3 mL) for 30 minutes at room temperature. The reaction mixture was added dropwise to ether (50 mL). The solution was allowed to stand at 4°C for 18 hours. The product was filtered, and dried under vacuum to yield 600 mg (79% yield) of the title compound as an off-white solid. Analysis calculated for C<sub>34</sub>H<sub>38</sub>N<sub>6</sub>O<sub>4</sub>S·1.3CF<sub>3</sub>CO<sub>2</sub>H: C, 56.72; H, 5.11; N, 10.84. Found: C, 56.34; H, 5.47; N, 11.49. The salt had a calculated molecular weight of 740.8.

Example 31

Preparation of N-[2-Benzylloxy-3-(N-benzylsulfonylamino)-6-methylphenyl-acetyl]-D,L-N- $\omega$ -4-methoxy-2,3,6-trimethylbenzenesulfonyl-D,L-3-amidinophenylalaninal-semicarbazonyl-4-N-diphenylmethane (42)

EDC (180.1 mg, 0.94 mmole) is added in one portion to a stirred solution of the compound of Example 8 (8) (209 mg, 0.49 mmole), hydroxybenzotriazole (75 mg, 0.49 mmole), and 4-methylmorpholine (0.24 mL, 2.2 mmole) in dimethylformamide (5

mL) with cooling in an ice bath. After 30 minutes, the compound of Example 30 (41) (363 mg, 0.49 mmole) is added. After an additional 22 hours, the reaction mixture is diluted with water (25 mL) and brine (25 mL). The product is  
5 filtered and dissolved in ethyl acetate (125 mL). The solution is washed with 10% citric acid, water, saturated sodium bicarbonate and brine, and is dried over anhydrous magnesium sulfate. The solvent is removed under vacuum. The resulting residue is chromatographed by flash chromatography  
10 on silica gel to give the title compound.

#### Example 32


Preparation of N-[2-Hydroxy-3-(N-benzyl-sulfonylamino)-6-methylphenyl-acetyl-D,L-3-amidinophenylalaninal semicarbazone  
15 (43)

The compound of Example 31 (42) (100 mg) is treated with hydrofluoric acid/anisole (9:1) for 30 minutes at -20°C and 0°C for 30 minutes. After removal of the hydrofluoric acid, the resulting residue is dissolved in 20% aqueous acetic acid  
20 and washed with diethyl ether. The aqueous layer is lyophilized to a powder, then is purified by preparative HPLC (C-18, eluting with 10-40% acetonitrile-water gradient containing 0.1% trifluoroacetic acid) to give the title compound.

25

#### Example 33

Preparation of N-[2-Hydroxy-3-(N-benzylsulfonylamino)-6-methylphenylacetyl]-D,L-3-amidinophenylalaninal (44)



30

The compound of Example 32 (43) (50 mg, 32 micromoles) is dissolved in methanol (1 mL) and 1% aqueous trifluoroacetic acid (5 mL), then formalin (0.23 mL) is added. After 40 minutes, the solution is filtered through a 2 micron filter, diluted to a volume of 15 mL with water, and then is purified by preparative HPLC (C-18, eluting with 10-40% acetonitrile-water gradient containing 0.1% trifluoroacetic acid). The fractions containing the title compound are pooled and lyophilized to give the title compound.

#### Example A

#### Kinetic Analysis of Compounds in an in vitro Thrombin Inhibition Assay

The ability of the compounds of Example 11 and 14 to act as an inhibitor of thrombin catalytic activity was assessed by determining the compounds' inhibition constant,  $K_i$ .

Enzyme activity was determined using the chromogenic substrate Pefachrome t-PA ( $\text{CH}_3\text{SO}_2$ -D-hexahydrotyrosine-glycyl-L-arginine-p-Nitroanilide), obtained from Pentapfarm Ltd. The substrate was reconstituted in deionized water prior to use. Purified human alpha-thrombin (3000U/mg specific activity) was obtained from Enzyme Research Laboratories, Inc. The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).

The assay for  $K_i$  determinations was conducted by combining in appropriate wells of a Corning microtiter plate, 100 microliters of HBSA or the test compound at a specified concentration diluted in HBSA (or HBSA alone for  $V_0$  (uninhibited velocity) measurement), and 100 microliters of the chromogenic substrate (750  $\mu\text{M}$ , 5X  $K_m$ ). At time zero, 50 microliters of alpha-thrombin diluted in HBSA were added to the wells, yielding a final concentration of 0.2 nM in a total volume of 250 microliters. Velocities of chromogenic substrate hydrolysis which occurred over 60 minutes were measured by the change in absorbance at 405 nm using a Thermo

Max<sup>®</sup> Kinetic Microplate Reader.  $K_i$  values were determined for test compounds using the relationships developed by Williams and Morrison, *Methods in Enzymology*, 63: 437 (1979) using steady state velocities ( $V_s$ ) measured over 60 minutes.

5 The extent of substrate hydrolysis was less than 5% over the course of this assay.

Table 1 below gives the  $K_i$  values for the compounds of Examples 11 and 14. The data show the utility of these

10 compounds as potent *in vitro* inhibitors of human alpha-thrombin.

Table 1

Compound	$K_i$ (nM)
Compound of Example 11	0.79
Compound of Example 14	1.0

15

#### Example B

##### In vitro Enzyme Assays for Specificity Determination

The ability of compounds of this invention to act as a

20 selective inhibitor of thrombin catalytic activity was assessed by determining the concentration of test compound which inhibited the activity of this enzyme by 50%, ( $IC_{50}$ ), and comparing this value to that determined for some or all of the following related serine proteases: recombinant tissue plasminogen activator (rt-PA), plasmin, activated protein C

25 (aPC), chymotrypsin and trypsin.

The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin). The assay for  $IC_{50}$  determinations was conducted by combining in

30 appropriate wells of a Corning microtiter plate, 50 microliters of HBSA, 50 microliters of the test compound at a specified concentration (covering a broad concentration



range) diluted in HBSA (or HBSA alone for  $V_0$  (uninhibited velocity) measurement), and 50 microliters of the enzyme diluted in HBSA. Following a 30-minute incubation at ambient temperature, 50 microliters of the substrate at the concentrations specified below were added to the wells, yielding a final total volume of 200 microliters. The initial velocity of chromogenic substrate hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max® Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was utilized. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC<sub>50</sub> value.

1. Thrombin (fIIa) Assay

Enzyme activity was determined using the chromogenic substrate, Pefachrome t-PA (CH<sub>3</sub>SO<sub>2</sub>-D-hexahydrotyrosine-glycyl-L-Arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was reconstituted in deionized water prior to use. Purified human  $\alpha$ -thrombin was obtained from Enzyme Research Laboratories, Inc. The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).

IC<sub>50</sub> determinations were conducted where HBSA (50  $\mu$ l),  $\alpha$ -thrombin (50  $\mu$ l) (the final enzyme concentration is 0.5 nM) and inhibitor (50  $\mu$ l) (covering a broad concentration range) were combined in appropriate wells and incubated for 30 minutes at room temperature prior to the addition of substrate Pefachrome-t-PA (50  $\mu$ l) (the final substrate concentration is 250  $\mu$ M, about 5 times  $K_m$ ). The initial velocity of Pefachrome t-PA hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max® Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was utilized. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC<sub>50</sub> value.

2. Recombinant tissue plasminogen activator (rt-PA)

rt-PA catalytic activity was determined using the substrate, Pefachrome t-PA ( $\text{CH}_3\text{SO}_2$ -D-hexahydrotyrosine-glycyl-L-arginine-p-nitroanilide, obtained from Pentapharm Ltd.). The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 500 micromolar (about 3-times  $K_m$ ). Human rt-PA (Activase®) was obtained from Genentech Inc. The enzyme was reconstituted in deionized water and diluted into HBSA prior to the assay in which the final concentration was 1.0 nM.

3. Plasmin Assay

Plasmin catalytic activity was determined using the chromogenic substrate, S-2366 (L-pyro-Glu-Pro-Arg-pNa·HCl), which was obtained from DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 300 micromolar (about 2.5-times  $K_m$ ). Purified human plasmin was obtained from Enzyme Research Laboratories, Inc. The enzyme was diluted into HBSA prior to assay in which the final concentration was 1.0 nM.

4. Activated Protein C (aPC)

aPC catalytic activity was determined using the chromogenic substrate, Pefachrome PC (delta-carbobenzloxy-D-lysine-L-prolyl-L-arginine-p-nitroanilide), obtained from Pentapharm Ltd.). The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 400 micromolar (about 3-times  $K_m$ ). Purified human aPC was obtained from Hematologic Technologies, Inc. The enzyme was diluted into HBSA prior to assay in which the final concentration was 1.0 nM.

5. Chymotrypsin

Chymotrypsin catalytic activity was determined using the chromogenic substrate, S-2586 (methoxy-succinyl-L-arginine-L-prolyl-L-tyrosyl-p-nitroanilide), which was obtained from  
5 DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 100 micromolar (about 9-times  $K_m$ ). Purified (3X-crystallized; CDI) bovine pancreatic  
10 a-chymotrypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

6. Trypsin

15 Trypsin catalytic activity was determined using the chromogenic substrate, S-2222 (benzoyl-L-isoleucine-L-glutamic acid-(gamma-methyl ester)-L-arginine-p-nitroanilide), which was obtained from DiaPharma Group. The  
20 substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 250 micromolar (about 4-times  $K_m$ ). Purified (3X-crystallized; TRL3) bovine pancreatic trypsin was obtained  
from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior  
25 to assay in which the final concentration was 0.5 nM.

7. Factor Xa

Factor Xa catalytic activity was determined using the chromogenic substrate S-2765 (N-benzyloxycarbonyl-D-arginine-L-glycine-L-arginine-p-nitroaniline), obtained from DiaPharma  
30 Group (Franklin, OH). All substrates were reconstituted in deionized water prior to use. The final concentration of S-2765 was 250  $\mu M$  (about 5-times  $K_m$ ). Purified human Factor X was obtained from Enzyme Research Laboratories, Inc. (South  
35 Bend, IN) and Factor Xa (FXa) was activated and prepared from it as described [Bock, P.E., Craig, P.A., Olson, S.T., and

Singh, P. Arch. Biochem. Biophys. 273:375-388 (1989)]. The enzyme was diluted into HBSA prior to assay, in which the final concentration was 0.25 nM.

5 Table 2 lists the determined IC<sub>50</sub> values for the compounds of Examples 11 and 14 as inhibitors of some of the enzymes listed above and demonstrates the high degree of specificity for the inhibition of alpha-thrombin compared to these related serine proteases.

10

Table 2  
IC<sub>50</sub> values for the inhibition of human alpha  
thrombin amidolytic activity compared to  
selected serine proteases for compounds of  
Example 11 and 14

15

Enzyme	Compound of Example 11	Compound of Example 14
	IC <sub>50</sub> (nM)	IC <sub>50</sub> (nM)
alpha-thrombin	3.19	1.44
rt-PA	NI*	NI*
Plasmin	NI*	NI*
aPC	NI*	NI*
Chymotrypsin	NI*	NI*
Trypsin	1145	1810

\*-No inhibition observed at the maximal concentration of inhibitor assayed-2500nM.

20

#### Example C

#### Ex Vivo Anticoagulant Effects of Compounds in Rat and Human Plasma

25 The ex vivo anticoagulant effects of compounds of the present invention are determined by measuring the prolongation of the activated partial thromboplastin time

(APTT) over a broad concentration range of the added inhibitor, using pooled normal human and rat plasma.

Fresh frozen citrated pooled normal human plasma is obtained from George King Biomedical, Overland Park, KA.

5 Pooled normal rat plasma is prepared from citrated whole blood collected from anesthetized rats using standard procedures. The plasma is flash frozen and stored at -80°C until use.

Measurements of APTT are made using the Coag-A-Mate RA4  
10 automated coagulometer (General Diagnostics, Organon Technica, Oklahoma City, OK) using the Automated APTT reagent (Organon Technica, Durham, NC) as the initiator of clotting according to the manufacturers instructions. The assay is conducted by making a series of dilutions of the test  
15 compounds into 100 µl rapidly thawed plasma followed by adding 200 microliters of APTT reagent and calcium chloride, to the wells of the assay carousel. Compounds that prolong the APTT in a dose-dependent manner are considered to demonstrate an anticoagulant effect.

20

#### Example D

#### Evaluation of the Antithrombotic Potential of Compounds in an Experimental Rat Model of Thrombosis

The antithrombotic (prevention of thrombus formation)  
25 properties of compounds of the present invention are evaluated using the following established experimental model of acute vascular thrombosis which is a rat model of FeCl<sub>3</sub>-induced platelet-dependent arterial thrombosis.

This is a well characterized model of platelet  
30 dependent, arterial thrombosis which has been used in the evaluation potential antithrombotic compounds such as direct thrombin inhibitors. See, Kurz, K. D., Main, B. W., and Sandusky, G. E., Thromb. Res., 60: 269-280 (1990). In this model a platelet-rich, occlusive thrombus is formed in a  
35 segment of the rat carotid artery treated locally with a fresh solution of FeCl<sub>3</sub> absorbed to a piece of filter paper.

The  $\text{FeCl}_3$  is thought to diffuse into the treated segment of artery and causes de-endothelialization of the affected vessel surface. This results in the exposure of blood to subendothelial structures which in turn causes platelet adherence, thrombin formation and platelet aggregation resulting in occlusive thrombus formation. The effect of a test compound on the incidence of occlusive thrombus formation following the application of the  $\text{FeCl}_3$  is monitored by ultrasonic flowtometry and is used as the primary end point. The use of flowtometry to measure carotid artery blood flow, is a modification of the original procedure in which thermal detection of clot formation was employed. See, Kurz, K. D., Main, B. W., and Sandusky, G. E., *Thromb. Res.*, 60: 269-280 (1990).

Male Harlan Sprague Dawley rats (420-450 g) are acclimated at least 72 hours prior to use and fasted for 12 hours prior to surgery with free access to water. The animals are prepared, anesthetized with Nembutal, followed by the insertion of catheters for blood pressure monitoring, drug and anesthesia delivery. The left carotid artery is isolated by making a midline cervical incision followed by blunt dissection and spreading techniques to separate a 2 cm segment of the vessel from the carotid sheath. A silk suture is inserted under the proximal and distal ends of the isolated vessel to provide clearance for the placement of a ultrasonic flow probe (Transonic) around the proximal end of the vessel. The probe is then secured with a stationary arm.

Following surgery the animals are randomized in either a control (saline) or treatment group with test compound, with at least 6 animals per group per dose. The test compound is administered as a single intravenous bolus at doses of about 0.3 to 5.0 mg/kg after placement of the flow probe and 5 minutes prior to the thrombogenic stimulus. At  $t=0$ , a 3mm diameter piece of filter paper (Whatman #3) soaked with 10 microliters of a 35% solution of fresh  $\text{FeCl}_3$  (made up in water) is applied the segment of isolated carotid artery,

distal to the flow probe. Blood pressure, blood flow, heart rate, and respiration are monitored for 60 minutes.

The incidence of occlusion (defined as the attainment of zero blood flow) is recorded as the primary end point. The efficacy of test compounds as an antithrombotic agent in preventing thrombus formation in this *in vivo* model is demonstrated by the reduction in the incidence of thrombotic occlusion. The effective dose which prevents 50% of thrombotic occlusions in this model (ED<sub>50</sub>) can be determined from the above data by plotting the incidence of occlusion versus the dose administered. This allows a direct comparison of the antithrombotic efficacy of test compounds with other antithrombotic agents which have also been evaluated in this model as . Table 3 lists the ED<sub>50</sub> values for several well known anticoagulant agents in this model (Heparin-Hirsh, J. N. Engl. J. Med., 324:1565-1574 (1992), Cairns, J.A. et. al. Chest, 102:456S-481S (1992); Argatroban-Gold, H.K. et.al., J. Am. Coll. Cardiol., 21:1039-1047 (1993); and Hirulog<sup>™</sup>-Sharma, G.V.R.K. et.al., Am. J. Cardiol., 72:1357-1360 (1993) and Lidón, R.M. et.al., Circulation, 88:1495-1501 (1993)).

Table 3

Efficacy of known antithrombotic agents based on ED<sub>50</sub> for thrombus prevention in the FeCl<sub>3</sub> model of arterial thrombosis

Compound	ED <sub>50</sub> <sup>a</sup>
Standard Heparin	200 U/kg
Argatroban	3.8 mg/kg
Hirulog <sup>™</sup>	3.0 mg/kg

<sup>a</sup>ED<sub>50</sub> is defined as the dose that prevents the incidence of complete thrombotic occlusion in 50% of animals tested.

The in vivo comparison of compounds of the present invention with the clinically effective antithrombotic agents Standard Heparin, Argatroban, and Hirulog<sup>tm</sup> in the same rodent model of experimental thrombosis coupled with  
5 demonstrated anticoagulant effects of the present compounds in both rat and human plasma described above in Example C would lead one skilled in the art to conclude that this compound will be an effective antithrombotic agent in humans.



Example EElastase Assay


Elastase catalytic activity was determined using the chromogenic substrate Elastase Substrate I (MeOSuc-Ala-Ala-Pro-Val-p-nitroaniline, obtained from Calbiochem). The substrate was reconstituted with 10% DMSO and assay buffer. Human Neutrophil Elastase was obtained from Calbiochem and diluted in the assay buffer prior to use. The assay buffer was diluted in the assay buffer prior to use. The assay buffer was HBS (10 mM HEPES, 150 mM sodium chloride, pH 7.5) with 0.1% Tween 80 and 10% DMSO. The DMSO is necessary to keep the substrate from precipitating during the course of the assay.

IC<sub>50</sub> determinations were conducted by combining in appropriate wells of a Corning microtiter plate, 50 microliters of the test compound at a specified concentration (covering a broad concentration range) diluted in buffer (or buffer along for V<sub>0</sub> (uninhibited velocity) measurement), and 50 microliters of the enzyme diluted in buffer, yielding an enzyme concentration of 1 nM. Following a 30 minute incubation at ambient temperature, 100 microliters of the substrate were added to the wells, yielding a final volume of 200 microliters, and a substrate concentration of 750  $\mu$ M (about 6 times K<sub>m</sub>).

The initial velocity of substrate hydrolysis was measured by the change of absorbance at 405 nm using a Thermo Max® Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was utilized. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC<sub>50</sub> value.

We claim:

1. A compound of the formula:



5

wherein

- (a) X is selected from the group consisting of  $-S(O)_2-$ ,  $-N(R')-S(O)_2-$ ,  $-C(=O)-$ ,  $-OC(=O)-$ ,  $-NH-C(=O)-$ ,  $-P(O)(R'')-$  and a direct link, wherein R' is hydrogen, alkyl of 1 to about 4 carbon atoms, aryl of about 6 to about 14 carbon atoms or aralkyl of about 6 to about 16 carbon atoms, and R'' is  $-NHR'$ ,  $-OR'$ ,  $-R'$ , or  $-SR'$ ;

- (b) R<sub>1</sub> is selected from the group consisting of:
- (1) alkyl of 1 to about 12 carbon atoms,
  - (2) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 5 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,



- (3) cycloalkyl of 3 to about 15 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,

- (4) heterocycloalkyl of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and  $S(O)_i$ , wherein i is 0, 1 or 2, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or  $-CO_2R'$ ,



- (5) heterocyclo f 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',
- (6) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',
- (7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,
- (8) heteroaryl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,
- (9) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,
- (10) heteroaralkyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,
- (11) aralkenyl of about 8 to about 16 carbon atoms having 6 to about 14 ring atoms which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,
- (12) heteroaralkenyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the

heteroatoms are selected from oxygen, nitrogen, and  $S(O)_i$ , and which is optionally mono-, di- or tri-substituted on the ring carbons with  $Y_1$ ,  $Y_2$ , and/or  $Y_3$ ,

5



10



15

(17) perfluoroalkyl of 1 to about 12 carbon atoms,

(18) perfluoroaryl of about 6 to about 14 carbon atoms,

(19) perfluoroaralkyl of about 7 to about 15 carbon atoms,

(20) hydrogen, and

20

(21) , wherein is a 5 to 7 membered heterocyclic ring having 3 to 6 ring carbon atoms, where V is  $-CH_2-$ ,  $-O-$ ,  $-S(=O)-$ ,  $-S(O)_2-$  or  $-S-$ , and wherein  $Y_1$ ,  $Y_2$ , and  $Y_3$  are

(i) independently selected from the group consisting of hydrogen, halogen, cyano, nitro, tetrazolyl,

25


amino, guanidin, amidino, methylamino, methylguanidin,  
 $-\text{CF}_3$ ,  $-\text{CF}_2\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{CH}(\text{CF}_3)_2$ ,  $-\text{C}(\text{OH})(\text{CF}_3)_2$ ,  $-\text{OCF}_3$ ,  
 $-\text{OCF}_2\text{CF}_3$ ,  $-\text{OC}(\text{O})\text{NH}_2$ ,  $-\text{OC}(\text{O})\text{NHZ}_1$ ,  $-\text{OC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{NHC}(\text{O})\text{Z}_1$ ,  
 $-\text{NHC}(\text{O})\text{NH}_2$ ,  $-\text{NHC}(\text{O})\text{NHZ}_1$ ,  $-\text{NHC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{NH}_2$ ,  
 5  $-\text{C}(\text{O})\text{NHZ}_1$ ,  $-\text{C}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OZ}_1$ ,  $-\text{P}(\text{O})_3\text{H}$ ,  $-\text{P}(\text{O})_3\text{H}_2$ ,  $-\text{PH}(\text{O})\text{OH}$ ,  
 $-\text{P}(\text{O})_3(\text{Z}_1)_2$ ,  $-\text{S}(\text{O})_3\text{H}$ ,  $-\text{S}(\text{O})_m\text{Z}_1$ ,  $-\text{Z}_1$ ,  $-\text{OZ}_1$ ,  $-\text{OH}$ ,  $-\text{NH}_2$ ,  $-\text{NHZ}_1$ ,  
 and  $-\text{NZ}_1\text{Z}_2$ , wherein  $m$  is 0, 1 or 2, and  $\text{Z}_1$  and  $\text{Z}_2$  are

independently selected from the group consisting of alkyl of  
 1 to about 12 carbon atoms, aryl of about 6 to about 14  
 10 carbon atoms, heteroaryl of about 5 to about 14 atoms having  
 1 to about 9 carbon atoms, aralkyl of about 7 to about 15  
 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms  
 having about 3 to about 9 carbon atoms, or

(ii)  $\text{Y}_1$  and  $\text{Y}_2$  are selected together to be  
 15  $-\text{OC}[(\text{Z}_3)(\text{Z}_4)]_q\text{O}-$ , wherein  $q$  is an integer from 1 to 4 and  $\text{Z}_3$   
 and  $\text{Z}_4$  are independently selected from the group consisting  
 of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of  
 about 6 to about 14 carbon atoms heteroaryl of about 5 to  
 about 14 atoms having 1 to about 9 carbon atoms, aralkyl of  
 20 about 7 to about 15 carbon atoms, and heteroaralkyl of about  
 6 to about 11 atoms having about 3 to about 9 carbon atoms,

(c)  $\text{R}_2$  is selected from the group consisting of  
 hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of  
 about 2 to about 4 carbon atoms,

25 (d)  $\text{R}_3$  is selected from the group consisting of



where  $d$  is an integer from 0 to 5 and  $\text{W}$  is  $-\text{N}-$  or  $-\text{CH}-$ ;

(e)  $\text{R}_4$  is selected from the group consisting of

(i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein  $n$  is 0, 1  
 30 or 2, and  $\text{R}_1$  is as defined above, with the proviso that  $\text{R}_1$  is  
 not a camphor derivative or  $-\text{N}(\text{Y})$  heterocycle,

- (ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ ,
- (iii) halogen,
- (iv) alkyl of 1 to 6 carbon atoms substituted with  $\text{Z}_5$ , wherein  $\text{Z}_5$  is selected from the group consisting of
  - 5 hydrogen,  $-\text{OH}$ ,  $-\text{OR}_8$ ,  $-\text{CH}_3$ , halogen,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{OR}_8$  and  $-\text{S}(\text{O})_p\text{R}_8$ , wherein  $\text{R}_8$  is alkyl of 1 to about 6 carbon atoms, and  $p$  is 0, 1 or 2,
- (v) alkyl of 1 to 3 carbon atoms substituted with cycloalkyl of 3 to 5 carbon atoms,
- 10 (vi) alkenyl of about 3 to about 6 carbon atoms,
- (vii) cycloalkyl of about 3 to about 10 carbon atoms,
- (viii) heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of
  - 15 the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, and
- (xi) heteroaralkyl of about 5 to about 10 ring atoms having 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms
  - 20 heteroatoms selected from oxygen, sulfur and nitrogen;
- (f)  $\text{R}_5$  is selected from the group consisting of
  - (i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ ,  $-\text{S}(\text{O})_n\text{R}_1$ , wherein  $\text{R}_1$  is as defined above, with the proviso that  $\text{R}_1$  is not a camphor derivative or  $\text{—}\overset{\text{N}}{\text{V}}$  heterocycle,
  - 25 (ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $\text{OCF}_2\text{H}$ , and
  - (iii) halogen;
- (g)  $\text{R}_6$  is selected from the group consisting of
  - (i)  $-\text{R}_1$ ,  $-\text{OR}_1$ ,  $-\text{NHR}_1$ , or  $-\text{S}(\text{O})_n\text{R}_1$ , wherein  $\text{R}_1$  is as defined above, with the proviso that  $\text{R}_1$  is not a camphor derivative or  $\text{—}\overset{\text{N}}{\text{V}}$  heterocycle,
  - 30 (ii)  $-\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $-\text{OCF}_3$ , or  $-\text{OCF}_2\text{H}$ ,
  - (iii) halogen,
  - (iv) alkyl of 1 to about 12 carbon atoms substituted with  $\text{Z}_6$ , wherein  $\text{Z}_6$  is selected from the group


- consisting of hydrogen, hydroxy, hal gen, -OR<sub>9</sub>, -NHR<sub>9</sub>, -C(O)OH, -C(O)OR<sub>9</sub>, and -S(O)<sub>p</sub>R<sub>9</sub>, wherein R<sub>9</sub> is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon atoms, aralkyl of about 7 to about 12 carbon atoms,
- 5 heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, or heteroaralkyl of about 5 to about 10
- 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,
- 15 (v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms,
- (vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 6 to about 10 carbon atoms or
- 20 heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms,
- (vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-
- 25 substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,
- (viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl
- 30 chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,
- (ix) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and
- 35 (x) heteroaralkenyl of about 5 to about 10 ring atoms having 1 to 9 ring carbon atoms and the remainder of

the ring at ms heter atoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>; and

(h) R<sub>7</sub> is selected from the group consisting of

5 (i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above with the

proviso that R<sub>1</sub> is not an camphor derivative or heterocycle,



(ii) -CF<sub>3</sub> or CF<sub>2</sub>H,

(iii) alkyl of 1 to about 10 carbon atoms,  
10 optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
-CONR<sub>10</sub>R<sub>11</sub>, wherein each of R<sub>10</sub> and R<sub>11</sub> is independnetly  
selected and is alkyl of 1 to about 4 carbon atoms or  
hydrogen,

(iv) alkenyl of about 3 to about 10 carbon atoms,  
15 optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>,  
-SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms,  
optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>,  
or -CONR<sub>10</sub>R<sub>11</sub>,

20 (vi) heteroaryl of about 5 to about 10 ring atoms  
having about 1 to about 9 ring carbon atoms, optionally  
substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
-CONR<sub>10</sub>R<sub>11</sub>, and

(vii) alkyl of 1 to about 3 carbon atoms  
25 substituted with heteroaryl of about 5 to about 10 ring atoms  
having about 1 to about 9 carbon atoms, optionally  
substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, or  
pharmaceutically acceptable salts thereof.

30 2. A compound according to claim 1, wherein X is  
selected from the group consisting of a direct link, -SO<sub>2</sub>-,  
and -NH-S(O)<sub>2</sub>-.

35 3. A compound according to claim 2, wherein X is  
-SO<sub>2</sub>-.



4. A compound according to claim 1, wherein R<sub>1</sub> is selected from the group consisting of alkyl, aralkyl, and aryl.

5

5. A compound according to claim 4, wherein R<sub>1</sub> is aralkyl selected from substituted benzyl and unsubstituted benzyl, or is aryl selected from substituted phenyl, unsubstituted phenyl, substituted naphthyl and unsubstituted naphthyl.

10

6. A compound according to claim 5, wherein R<sub>1</sub> is aryl substituted with a group selected from the group consisting of methyl, methoxy, fluoro, chloro, trifluoromethyl, and -OCF<sub>3</sub>.

15

7. A compound according to claim 5, wherein R<sub>1</sub> is cyclohexyl or cyclohexylmethyl.


20

8. A compound according to claim 4, wherein R<sub>1</sub> is aralkyl.

9. A compound according to claim 1, wherein R<sub>2</sub> is hydrogen.

25

10. A compound according to claim 1, wherein R<sub>3</sub> is



11. A compound according to claim 1, wherein R<sub>4</sub> is selected from the group consisting of:  
(a) hydrogen,

30

(b) alkyl of 1 to 6 carbon atoms substituted with Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of hydrogen, -OH, -OR<sub>g</sub>, -CH<sub>3</sub>, halogen, -C(O)OH, -C(O)OR<sub>g</sub>, -S(O)<sub>3</sub>OH and -S(O)<sub>p</sub>R<sub>g</sub>.

5 (c) alkyl of 1 to 3 carbon atoms substituted with cyclic alkyl of 3 to 5 carbon atoms,

(d) alkenyl,

(e) cycloalkyl,

(f) heteroaryl,

10 (g) heteroaralkyl, and

(h) trifluoromethyl.

12. A compound according to claim 11, wherein R<sub>4</sub> is selected from the group consisting of methyl, ethyl,  
15 propenyl, allyl, propyl, isopropyl, butyl, (R)-sec-butyl, (S)-sec-butyl, isobutyl, 1-pentyl, (R)-2-pentyl, (S)-2-pentyl, 3-pentyl, (S)-1-(2-methyl)-butyl, (R)-2-(3-methyl)-butyl, 1-(3-methyl)-butyl, (R)-1-(2-methyl)-butyl, cyclopentyl, 2-pyrrolyl, 3-pyrrolyl, 1-hexyl, (S)-2-hexyl,  
20 (R)-2-hexyl, (R)-3-hexyl, (S)-3-hexyl, hydroxy, -OR<sub>g</sub>, wherein R<sub>g</sub> is alkyl of 1 to about 4 carbon atoms, phenyloxy, -OCF<sub>3</sub>, and halogen.

13. A compound according to claim 1, wherein R<sub>5</sub> is  
25 selected from the group consisting of hydrogen, halogen, alkyl of 1 to about 5 carbon atoms, trifluoromethyl, alkoxy of 1 to about 4 carbon atoms.

14. A compound according to claim 13, wherein R<sub>5</sub> is  
30 hydrogen.

15. A compound according to claim 1, wherein R<sub>6</sub> is selected from the group consisting of:

(a) hydrogen,

35 (b) alkyl of 1 to about 12 carbon atoms substituted with Z<sub>6</sub>.

(c) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl,

(d) alkenyl of about 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl,

5 (e) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri- substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

10 (f) heteroaralkyl ring of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

15 (g) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and,

20 (h) heteroaralkenyl of about 5 to about 10 ring atoms having 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>.

25 16. A compound according to claim 1, wherein R<sub>4</sub> and R<sub>5</sub> are independently selected from the group consisting of hydrogen and methyl and R<sub>6</sub> is selected from the group consisting of aralkyl of about 8 to about 13 carbon atoms, -O-aralkyl, -NH-aralkyl, and -S(O)<sub>p</sub>-aralkyl of about 7 to about 12 carbon atoms.

30

17. A compound according to claim 16, wherein said aralkyl group has an aryl group selected from unsubstituted phenyl, substituted phenyl, substituted naphthyl and unsubstituted naphthyl.

35

18. A compound according to claim 17, wherein said aryl

group is substituted with a substituent selected from the group consisting of methyl, methoxy, fluoro, chloro and trifluoromethyl.

5           19. A compound according to claim 15, wherein R<sub>6</sub> is selected from the group consisting of phenylethyl, phenylpropyl, cyclohexylethyl and cyclohexylpropyl.

10           20. A compound according to claim 1, wherein R<sub>7</sub> is selected from the group consisting of hydrogen, methyl, difluoromethyl and trifluoromethyl.

15           21. A compound according to claim 20, wherein R<sub>7</sub> is hydrogen or methyl.

            22. A compound according to claim 1, wherein R<sub>7</sub> is selected from the group consisting of:

(a) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

20           (b) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(c) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
25           -CONR<sub>10</sub>R<sub>11</sub>,

(d) heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 ring carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
-CONR<sub>10</sub>R<sub>11</sub>, and

30           (e) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, and -CONR<sub>10</sub>R<sub>11</sub>.


35           23. A compound according to claim 1, wherein R<sub>5</sub> and R<sub>6</sub> are independently selected from hydrogen and methyl, and R<sub>4</sub>

is selected from the group consisting of hydrogen, methyl, ethyl, propenyl, allyl, propyl, isopr pyl, butyl, (R)-sec-butyl, (S)-sec-butyl, isobutyl, 1-pentyl, (R)-2-pentyl, (S)-2-pentyl, 3-pentyl, (S)-1-(2-methyl)-butyl, (R)-2-(3-methyl)-butyl, 1-(3-methyl)-butyl, (R)-1-(2-methyl)-butyl, cyclopentyl, 1-hexyl, (S)-2-hexyl, (R)-2-hexyl, (R)-3-hexyl, and (S)-3-hexyl.

10 24. A compound according to claim 1, wherein R<sub>4</sub>, R<sub>5</sub>, and R<sub>6</sub>, are independently selected from hydrogen and methyl.

25. A compound according to claim 1, wherein any two of R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub>, and R<sub>6</sub> are hydrogen.


15 26. A compound according to claim 1, wherein X is -S(O)<sub>2</sub>-, R<sub>1</sub> is substituted or unsubstituted aralkyl, and R<sub>3</sub> is



20

27. A compound according to claim 26, wherein R<sub>1</sub> is substituted or unsubstituted benzyl.

28. A compound of the formula:



(II)

wherein,

- 5 (a) X is selected from the group consisting of  $-S(O)_2-$ ,  $-N(R')-S(O)_2-$ ,  $-(C=O)-$ ,  $-OC(=O)-$ ,  $-NH-C(=O)-$ ,  $-P(O)(R'')$  and a direct link, wherein  $R'$  is hydrogen, alkyl of 1 to about 4 carbon atoms, aryl of about 6 to about 14 carbon atoms or aralkyl of about 6 to about 16 carbon atoms, and  $R''$  is  $-NR'$ ,  $-OR'$ ,  $-R'$ , or  $-SR'$ ;
- 10 (b)  $R_1$  is selected from the group consisting of:
  - (1) alkyl of 1 to about 12 carbon atoms,
  - (2) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 5 to about 8 carbon atoms, which
    - 15 optionally is substituted in the ring carbons with hydroxyl, amino, guanidino, amidino,  $C_1$  to  $C_3$  alkyl,  $C_1$  to  $C_3$  alkoxy, or  $CO_2R'$ ,
  - (3) cycloalkyl of 3 to about 15 carbon atoms, which optionally is substituted in the ring carbons with
    - 20 hydroxyl, amino, guanidino, amidino,  $C_1$  to  $C_3$  alkyl,  $C_1$  to  $C_3$  alkoxy, or  $CO_2R'$ ,
  - (4) heterocycloalkyl of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group
    - 25 consisting of oxygen, nitrogen, and  $S(O)_i$ , wherein  $i$  is 0, 1 or 2, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino,  $C_1$  to  $C_3$  alkyl,  $C_1$

to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',

(5) heterocyclo of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',

(6) alkenyl of about 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',

(7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,





(8) heteroaryl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(9) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(10) heteroaralkyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,

(11) aralkenyl of about 8 to about 16 carbon atoms having 6 to about 14 ring atoms which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,

- (12) hetero aralkenyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,




15

- (17) perfluoroalkyl of 1 to about 12 carbon atoms,  
 (18) perfluoroaryl of about 6 to about 14 carbon atoms,  
 (19) perfluoroaryl alkyl of about 7 to about 15 carbon atoms,  
 (20) hydrogen, and

- (21)   
 , wherein is a 5 to 7 member heterocyclic ring having 3 to 6 ring carbon atoms, where V is -CH<sub>2</sub>-, -O-, -S(=O)-, -S(O)<sub>2</sub>- or -S-,  
 wherein Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>3</sub> are

25



- (i) independently selected from the group consisting of hydrogen, halogen, cyano, nitro, tetrazolyl, amino, guanidino, amidino, methylamino, methylguanidino, -CF<sub>3</sub>, -CF<sub>2</sub>CF<sub>3</sub>, -CF<sub>2</sub>H, CH(CF<sub>3</sub>)<sub>2</sub>, -C(OH)(CF<sub>3</sub>)<sub>2</sub>, -OCF<sub>3</sub>, -OCF<sub>2</sub>CF<sub>3</sub>, -OC(O)NH<sub>2</sub>, -OC(O)NHZ<sub>1</sub>, -OC(O)NZ<sub>1</sub>Z<sub>2</sub>, -NHC(O)Z<sub>1</sub>, -NHC(O)NH<sub>2</sub>, -NHC(O)NZ<sub>1</sub>, -NHC(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OH, -C(O)NH<sub>2</sub>, -C(O)NHZ<sub>1</sub>, -C(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OZ<sub>1</sub>, -P(O)<sub>3</sub>H, -P(O)<sub>3</sub>H<sub>2</sub>, -PH(O)OH, -P(O)<sub>3</sub>(Z<sub>1</sub>)<sub>2</sub>, -S(O)<sub>3</sub>H, -S(O)<sub>m</sub>Z<sub>1</sub>, -Z<sub>1</sub>, -OZ<sub>1</sub>, -OH, -NH<sub>2</sub>, -NHZ<sub>1</sub>, and -NZ<sub>1</sub>Z<sub>2</sub>, wherein m is 0, 1 or 2, and Z<sub>1</sub> and Z<sub>2</sub> are independently selected from the group consisting of alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 14 carbon atoms, heteroaryl of about 5 to about 14 atoms having 1 to about 9 carbon atoms, aralkyl of about 7 to about 15 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms having about 3 to about 9 carbon atoms, or
- (ii) Y<sub>1</sub> and Y<sub>2</sub> are selected together to be -OC[(Z<sub>3</sub>)(Z<sub>4</sub>)]<sub>q</sub>O-, wherein q is an integer from 1 to 4 and Z<sub>3</sub> and Z<sub>4</sub> are independently selected from the group consisting of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 14 carbon atoms, heteroaryl of about 5 to about 14 atoms having 1 to about 9 carbon atoms, aralkyl of about 7 to about 15 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms having about 3 to about 9 carbon atoms;
- (c) R<sub>2</sub> is selected from the group consisting of hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of about 2 to about 4 carbon atoms;
- (d) R<sub>4</sub> is selected from the group consisting of
- (i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein n is 0, 1 or 2, and R<sub>1</sub> is as defined above, with the proviso that R<sub>1</sub> is not a camphor derivative or  heterocycle,
- (ii) -CF<sub>3</sub>, -CF<sub>2</sub>H, -OCF<sub>3</sub>, -OCF<sub>2</sub>H,
- (iii) halogen,
- (iv) alkyl of 1 to 6 carbon atoms substituted with Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of hydrogen, -OH, -OR<sub>g</sub>, -CH<sub>3</sub>, halogen, -C(O)OH, -C(O)OR<sub>g</sub> and

-S(O)<sub>p</sub>R<sub>8</sub>, wherein R<sub>8</sub> is alkyl of 1 to about 6 carbon atoms, and p is 0, 1 or 2,

(v) alkyl of 1 to 3 carbon atoms substituted with cyclic alkyl of 3 to 5 carbon atoms,

5 (vi) alkenyl of about 3 to about 6 carbon atoms,


(vii) cycloalkyl of about 3 to about 10 carbon atoms,

(viii) heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of  
10 the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, and

(ix) heteroaralkyl of about 5 to about 10 ring atoms having 2 to about 15 carbon atoms which include 1 to  
15 about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen;

(e) R<sub>5</sub> is selected from the group consisting of

(i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein R<sub>1</sub> is as defined above, with the proviso that R<sub>1</sub> is not a camphor


derivative or  heterocycle,

20 (ii) -CF<sub>3</sub>, -CF<sub>2</sub>H, -OCF<sub>3</sub>, -OCF<sub>2</sub>H, and

(iii) halogen;

(f) R<sub>6</sub> is selected from the group consisting of

(i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein R<sub>1</sub> is as defined above, with the proviso that R<sub>1</sub> is not a camphor

25 derivative or  heterocycle,

(ii) -CF<sub>3</sub>, -CF<sub>2</sub>H, -OCF<sub>3</sub>, -OCF<sub>2</sub>H,

(iii) halogen,

(iv) alkyl of 1 to about 12 carbon atoms substituted with Z<sub>6</sub>, wherein Z<sub>6</sub> is selected from the group  
30 consisting of hydrogen, hydroxy, halogen, -OR<sub>9</sub>, -NHR<sub>9</sub>, -C(O)OH, -C(O)OR<sub>9</sub>, and -S(O)<sub>p</sub>R<sub>9</sub>, wherein R<sub>9</sub> is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon atoms, aralkyl of about 7 to about 12 carbon atoms, heteroaryl of about 5 to about 10 ring atoms 1 to about 9

ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted in the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and heteroaralkyl of about 5

5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted in the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

10 (v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms,

(vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 6 to about 10 carbon atoms, or  
15 heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms,

(vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-  
20 substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl  
25 chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,


(ix) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

30 (x) heteroaralkenyl ring of about 5 to about 10 ring atoms with 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>;

35 (g) R<sub>7</sub> is selected from the group consisting of

(i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above, with the

pr viso that R<sub>7</sub> is not an R<sub>1</sub> camphor derivative or heterocycle,



(ii) -CF<sub>3</sub> or CF<sub>2</sub>H,

(iii) alkyl of 1 to about 10 carbon atoms,  
5 optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>,  
-CONR<sub>10</sub>R<sub>11</sub>, wherein each of R<sub>1</sub> and R<sub>11</sub> is independently  
selected and is alkyl of 1 to about 4 carbon atoms or  
hydrogen,

(iv) alkenyl of about 3 to about 10 carbon atoms,  
10 optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>,  
CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms,  
optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>,  
-CONR<sub>10</sub>R<sub>11</sub>,

15 (vi) heteroaryl of about 5 to about 10 ring atoms,  
having about 1 to about 9 ring carbon atoms which is  
optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
-CONR<sub>10</sub>R<sub>11</sub>, and

(vii) alkyl of 1 to about 3 carbon atoms  
20 substituted with heteroaryl of about 5 to about 16 atoms  
having 1 to about 9 carbon atoms, which is optionally  
substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, CONR<sub>10</sub>R<sub>11</sub>; and

(h) R<sub>12</sub> is selected from the group consisting of

(i) alkyl of 1 to about 12 carbon atoms which is  
25 optionally substituted with amino, amidino or guanidino,

(ii) aralkyl of 7 to about 15 carbon atoms which is  
optionally substituted on a ring carbon with amino, amidino,  
or guanidino,

(iii) aryl of about 6 to about 14 carbon atoms which  
30 is optionally substituted on a ring carbon with amino,  
amidino, or guanidino,


(iv) heteroaralkyl of 5 to about 14 ring atoms  
having 1 to about 9 ring carbon atoms and the remainder of  
the ring atoms heteroatoms selected from oxygen, nitrogen and  
35 sulfur which is optionally substituted on a ring carbon with

amino, amidino or guanidino; and

- (j) TS is selected from the group consisting of
- (i) -CHO,
  - (ii) -C(O)CF<sub>3</sub>,
  - 5 (iii) -C(O)CF<sub>2</sub>CF<sub>3</sub>,
  - (iv) -C(O)-C(O)-NHR<sub>13</sub>,
  - (v) -C(O)-C(O)<sub>2</sub>R<sub>13</sub>,
  - (vi) -C(O)-C(O)-NR<sub>13</sub>R<sub>14</sub>,
  - (vii) -B(OR<sub>13</sub>)(OR<sub>14</sub>),
  - 10 (viii) -C(O)CH<sub>2</sub>Cl, and
  - (ix) -C(O)-R<sub>15</sub>;

wherein R<sub>13</sub> and R<sub>14</sub> are independently selected from hydrogen  
lower alkyl of 1 to about 6 carbon atoms, cycloalkyl of 3 to  
about 10 carbon atoms, aralkyl of 7 to about 15 carbon atoms,  
15 and heteroaralkyl of about 5 to about 10 ring atoms having  
about 2 to about 15 carbon atoms which include 1 to about 9  
ring carbon atoms and the remainder of the ring atoms  
heteroatoms selected from oxygen, nitrogen and sulfur; or  
when TS is -B(OR<sub>13</sub>)(OR<sub>14</sub>) then OR<sub>13</sub> and OR<sub>14</sub> may be taken  
20 together to be -O[C(Z<sub>3</sub>)(Z<sub>4</sub>)]qO-; and R<sub>15</sub> is selected from aryl  
of 6 to about 14 carbon atoms and heteroaryl of 5 to about 14  
ring atoms having 1 to about 13 ring carbon atoms and the  
remainder of the ring atoms selected from oxygen, nitrogen  
and sulfur; and pharmaceutically acceptable salts thereof.

29. A compound of the formula:



5 wherein

(a) X is selected from the group consisting of -S(O)<sub>2</sub>-,  
 -N(R')-S(O)<sub>2</sub>-, -C(=O)-, -OC(=O)-, -NH-C(=O)-, -P(O)(R'')- and  
 a direct link, wherein R' is hydrogen, alkyl of 1 to about 4  
 carbon atoms, aryl of about 6 to about 14 carbon atoms or  
 10 aralkyl of about 6 to about 16 carbon atoms, and R'' is -NHR',  
 -OR', -R', or -SR';

(b) R<sub>1</sub> is selected from the group consisting of:

(1) alkyl of 1 to about 12 carbon atoms,  
 (2) alkyl of 1 to about 3 carbon atoms substituted  
 15 with cycloalkyl of about 5 to about 8 carbon atoms, which  
 optionally is substituted on the ring carbons with hydroxyl,  
 amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy,  
 or -CO<sub>2</sub>R',

(3) cycloalkyl of 3 to about 15 carbon atoms,  
 20 which optionally is substituted on the ring carbons with  
 hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub>  
 alkoxy, or -CO<sub>2</sub>R',

(4) heterocycloalkyl of 4 to about 10 ring atoms  
 with the ring atoms selected from carbon and heteroatoms,  
 25 wherein the heteroatoms are selected from the group  
 consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, wherein i is 0, 1  
 or 2, and which is optionally substituted on the ring carbons  
 with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub>  
 to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

30 (5) heterocyclo of 4 to about 10 ring atoms with

the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally substituted on the ring carbons with hydroxyl, amino, 5 guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

(6) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, which optionally is substituted on the ring 10 carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or -CO<sub>2</sub>R',

(7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

15 (8) heteroaryl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,





20 (9) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(10) heteroaralkyl of 5 to 14 ring atoms with the 25 ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

30 (11) aralkenyl of about 8 to about 16 carbon atoms having 6 to about 14 ring atoms which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(12) heteroaralkenyl of 5 to 14 ring atoms with the 35 ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>,

and which is opti nally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,





(17) perfluoroalkyl of 1 to about 12 carbon atoms,

(18) perfluoroaryl of about 6 to about 14 carbon

15 atoms,

(19) perfluoroaralkyl of about 7 to about 15 carbon atoms,

(20) hydrogen, and

(21) , wherein  is a 5 to 7 membered heterocyclic ring having 3 to 6 ring carbon atoms, where V is -CH<sub>2</sub>-, -O-, -S(=O)-, -S(O)<sub>2</sub>- or -S-, and wherein Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>3</sub> are

(i) independently selected from the group consisting of hydrogen, halogen, cyano, nitro, tetrazolyl, amino, guanidino, amidino, methylamino, methylguanidino,

25



- CF<sub>3</sub>, -CF<sub>2</sub>CF<sub>3</sub>, -CF<sub>2</sub>H, -CH(CF<sub>3</sub>)<sub>2</sub>, -C(OH)(CF<sub>3</sub>)<sub>2</sub>, -OCF<sub>3</sub>,  
 -OCF<sub>2</sub>CF<sub>3</sub>, -OC(O)NH<sub>2</sub>, -OC(O)NHZ<sub>1</sub>, -OC(O)NZ<sub>1</sub>Z<sub>2</sub>, -NHC(O)Z<sub>1</sub>,  
 -NHC(O)NH<sub>2</sub>, -NHC(O)NHZ<sub>1</sub>, -NHC(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OH, -C(O)NH<sub>2</sub>,  
 -C(O)NHZ<sub>1</sub>, -C(O)NZ<sub>1</sub>Z<sub>2</sub>, -C(O)OZ<sub>1</sub>, -P(O)<sub>3</sub>H, -P(O)<sub>3</sub>H<sub>2</sub>, -PH(O)OH,  
 5 -P(O)<sub>3</sub>(Z<sub>1</sub>)<sub>2</sub>, -S(O)<sub>3</sub>H, -S(O)<sub>m</sub>Z<sub>1</sub>, -Z<sub>1</sub>, -OZ<sub>1</sub>, -OH, -NH<sub>2</sub>, -NHZ<sub>1</sub>,  
 and -NZ<sub>1</sub>Z<sub>2</sub>, wherein m is 0, 1 or 2, and Z<sub>1</sub> and Z<sub>2</sub> are


independently selected from the group consisting of alkyl of  
 1 to about 12 carbon atoms, aryl of about 6 to about 14  
 carbon atoms, heteroaryl of about 5 to about 14 atoms having  
 10 1 to about 9 carbon atoms, aralkyl of about 7 to about 15  
 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms  
 having about 3 to about 9 carbon atoms, or

- (ii) Y<sub>1</sub> and Y<sub>2</sub> are selected together to be  
 -OC[(Z<sub>3</sub>)(Z<sub>4</sub>)]<sub>q</sub>O-, wherein q is an integer from 1 to 4 and Z<sub>3</sub>  
 15 and Z<sub>4</sub> are independently selected from the group consisting  
 of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of  
 about 6 to about 14 carbon atoms heteroaryl of about 5 to  
 about 14 atoms having 1 to about 9 carbon atoms, aralkyl of  
 about 7 to about 15 carbon atoms, and heteroaralkyl of about  
 20 6 to about 11 atoms having about 3 to about 9 carbon atoms,

(c) R<sub>2</sub> is selected from the group consisting of  
 hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of  
 about 2 to about 4 carbon atoms,

(d) R<sub>3</sub> is

25



, where d is an integer from 0 to 5;

(e) R<sub>4</sub> is selected from the group consisting of

- (i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein n is 0, 1  
 30 or 2, and R<sub>1</sub> is as defined above, with the proviso that R<sub>1</sub> is

not a camphor derivative or  $\text{---}\overset{\text{N}}{\text{---}}\text{---}\overset{\text{Y}}{\text{---}}$  heterocycle,

(ii)  $\text{---CF}_3$ ,  $\text{---CF}_2\text{H}$ ,  $\text{---OCF}_3$ , or  $\text{---OCF}_2\text{H}$ ,

(iii) halogen,

(iv) alkyl of 1 to 6 carbon atoms substituted with  
5 Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of  
hydrogen,  $\text{---OH}$ ,  $\text{---OR}_8$ ,  $\text{---CH}_3$ , halogen,  $\text{---C(O)OH}$ ,  $\text{---C(O)OR}_8$  and  
 $\text{---S(O)}_p\text{R}_8$ , wherein R<sub>8</sub> is alkyl of 1 to about 6 carbon atoms,  
and p is 0, 1 or 2,

(v) alkyl of 1 to 3 carbon atoms substituted with  
10 cycloalkyl of 3 to 5 carbon atoms,

(vi) alkenyl of about 3 to about 6 carbon atoms,

(vii) cycloalkyl of about 3 to about 10 carbon  
atoms,

(viii) heteroaryl of about 5 to about 10 ring atoms  
15 having 1 to about 9 ring carbon atoms and the remainder of  
the ring atoms heteroatoms selected from oxygen, sulfur and  
nitrogen, and

(xi) heteroaralkyl of about 5 to about 10 ring  
atoms having 2 to about 15 carbon atoms which include 1 to  
20 about 9 ring carbon atoms and the remainder of the ring atoms  
heteroatoms selected from oxygen, sulfur and nitrogen;

(f) R<sub>5</sub> is selected from the group consisting of

(i)  $\text{---R}_1$ ,  $\text{---OR}_1$ ,  $\text{---NHR}_1$ ,  $\text{---S(O)}_n\text{R}_1$ , wherein R<sub>1</sub> is as  
defined above, with the proviso that R<sub>1</sub> is not a camphor

25 derivative or  $\text{---}\overset{\text{N}}{\text{---}}\text{---}\overset{\text{Y}}{\text{---}}$  heterocycle,

(ii)  $\text{---CF}_3$ ,  $\text{---CF}_2\text{H}$ ,  $\text{---OCF}_3$ , or  $\text{---OCF}_2\text{H}$ , and

(iii) halogen;

(g) R<sub>6</sub> is selected from the group consisting of

(i)  $\text{---R}_1$ ,  $\text{---OR}_1$ ,  $\text{---NHR}_1$ , or  $\text{---S(O)}_n\text{R}_1$ , wherein R<sub>1</sub> is as  
30 defined above, with the proviso that R<sub>1</sub> is not a camphor

derivative or  $\text{---}\overset{\text{N}}{\text{---}}\text{---}\overset{\text{Y}}{\text{---}}$  heterocycle,

(ii)  $\text{---CF}_3$ ,  $\text{---CF}_2\text{H}$ ,  $\text{---OCF}_3$ , or  $\text{---OCF}_2\text{H}$ ,

(iii) halogen,

(iv) alkyl of 1 to about 12 carbon atoms

substituted with Z<sub>6</sub>, wherein Z<sub>6</sub> is selected from the group consisting of hydrogen, hydroxy, halogen, -OR<sub>9</sub>, -NHR<sub>9</sub>, -C(O)OH, -C(O)OR<sub>9</sub>, and -S(O)<sub>p</sub>R<sub>9</sub>, wherein R<sub>9</sub> is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon atoms, aralkyl of about 7 to about 12 carbon atoms, heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, or heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms,

(vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, aryl of about 6 to about 10 carbon atoms or heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms,

(vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri- substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(ix) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

(x) heteroaralkenyl of about 5 to about 10 ring

atoms having 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>; and

5 (h) R<sub>7</sub> is selected from the group consisting of

(i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above with the

proviso that R<sub>1</sub> is not an camphor derivative or heterocycle,



(ii) -CF<sub>3</sub> or CF<sub>2</sub>H,

10 (iii) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, wherein each of R<sub>10</sub> and R<sub>11</sub> is independently selected and is alkyl of 1 to about 4 carbon atoms or hydrogen,

15 (iv) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

20 (vi) heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 ring carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, and

25 (vii) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>, or pharmaceutically acceptable salts thereof.

30

30. A compound according to claim 29, wherein X is selected from the group consisting of a direct link, -SO<sub>2</sub>-, and -NH-S(O)<sub>2</sub>-.

35 31. A compound according to claim 30, wherein X is

-SO<sub>2</sub>-.

32. A compound according to claim 29, wherein R<sub>1</sub> is selected from the group consisting of alkyl, aralkyl, and aryl.

33. A compound according to claim 32, wherein R<sub>1</sub> is aralkyl selected from substituted benzyl and unsubstituted benzyl, or is aryl selected from substituted phenyl, unsubstituted phenyl, substituted naphthyl and unsubstituted naphthyl.

34. A compound according to claim 33, wherein R<sub>1</sub> is aryl substituted with a group selected from the group consisting of methyl, methoxy, fluoro, chloro, trifluoromethyl, and -OCF<sub>3</sub>.

35. A compound according to claim 33, wherein R<sub>1</sub> is cyclohexyl or cyclohexylmethyl.

36. A compound according to claim 32, wherein R<sub>1</sub> is aralkyl.

37. A compound according to claim 29, wherein R<sub>2</sub> is hydrogen.

38. A compound according to claim 29, wherein R<sub>4</sub> is selected from the group consisting of:

- (a) hydrogen,
- (b) alkyl of 1 to 6 carbon atoms substituted with Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of hydrogen, -OH, -OR<sub>8</sub>, -CH<sub>3</sub>, halogen, -C(O)OH, -C(O)OR<sub>8</sub>, -S(O)<sub>3</sub>OH and -S(O)<sub>p</sub>R<sub>8</sub>,
- (c) alkyl of 1 to 3 carbon atoms substituted with cyclic alkyl of 3 to 5 carbon atoms,
- (d) alkenyl,

- (e) cycloalkyl,
- (f) heteroaryl,
- (g) heteroaralkyl, and
- (h) trifluoromethyl.

5

39. A compound according to claim 38, wherein R<sub>4</sub> is selected from the group consisting of methyl, ethyl, propenyl, allyl, propyl, isopropyl, butyl, (R)-sec-butyl, (S)-sec-butyl, isobutyl, 1-pentyl, (R)-2-pentyl, (S)-2-pentyl, 3-pentyl, (S)-1-(2-methyl)-butyl, (R)-2-(3-methyl)-butyl, 1-(3-methyl)-butyl, (R)-1-(2-methyl)-butyl, cyclopentyl, 2-pyrrolyl, 3-pyrrolyl, 1-hexyl, (S)-2-hexyl, (R)-2-hexyl, (R)-3-hexyl, (S)-3-hexyl, hydroxy, -OR<sub>8</sub>, wherein R<sub>8</sub> is alkyl of 1 to about 4 carbon atoms, phenyloxy, -OCF<sub>3</sub>, and halogen.

15<sup>f</sup>

40. A compound according to claim 29, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, halogen, alkyl of 1 to about 5 carbon atoms, trifluoromethyl, alkoxy of 1 to about 4 carbon atoms.

20

41. A compound according to claim 40, wherein R<sub>5</sub> is hydrogen.

25

42. A compound according to claim 29, wherein R<sub>6</sub> is selected from the group consisting of:

- (a) hydrogen,
- (b) alkyl of 1 to about 12 carbon atoms substituted with Z<sub>6</sub>,

26,

30

(c) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl,

(d) alkenyl of about 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl,

(e) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri- substituted with

35

Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(f) heteroaralkyl ring of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(g) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and,

(h) heteroaralkenyl of about 5 to about 10 ring atoms having 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>.

43. A compound according to claim 29, wherein R<sub>4</sub> and R<sub>5</sub> are independently selected from the group consisting of hydrogen and methyl and R<sub>6</sub> is selected from the group consisting of aralkyl of about 8 to about 13 carbon atoms, -O-aralkyl, -NH-aralkyl, and -S(O)<sub>p</sub>-aralkyl of about 7 to about 12 carbon atoms.

44. A compound according to claim 43, wherein said aralkyl group has an aryl group selected from unsubstituted phenyl, substituted phenyl, substituted naphthyl and unsubstituted naphthyl.

45. A compound according to claim 44, wherein said aryl group is substituted with a substituent selected from the group consisting of methyl, methoxy, fluoro, chloro and trifluoromethyl.

46. A compound according to claim 42, wherein R<sub>6</sub> is selected from the group consisting of phenylethyl, phenylpropyl, cyclohexylethyl and cyclohexylpropyl.

47. A compound according to claim 29, wherein R<sub>7</sub> is selected from the group consisting of hydrogen, methyl, difluoromethyl and trifluoromethyl.

5

48. A compound according to claim 47, wherein R<sub>7</sub> is hydrogen or methyl.

49. A compound according to claim 29, wherein R<sub>7</sub> is selected from the group consisting of:

(a) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(b) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>,  
15 for -CONR<sub>10</sub>R<sub>11</sub>,

(c) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or -CONR<sub>10</sub>R<sub>11</sub>,

(d) heteroaryl of about 5 to about 10 ring atoms having about 1 to about 9 ring carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or  
20 -CONR<sub>10</sub>R<sub>11</sub>, and

(e) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 10 ring atoms having about 1  
25 to about 9 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, and -CONR<sub>10</sub>R<sub>11</sub>.

50. A compound according to claim 29, wherein R<sub>5</sub> and R<sub>6</sub> are independently selected from hydrogen and methyl, and R<sub>4</sub>  
30 is selected from the group consisting of hydrogen, methyl, ethyl, propenyl, allyl, propyl, isopropyl, butyl, (R)-sec-butyl, (S)-sec-butyl, isobutyl, 1-pentyl, (R)-2-pentyl, (S)-2-pentyl, 3-pentyl, (S)-1-(2-methyl)-butyl, (R)-2-(3-methyl)-butyl, 1-(3-methyl)-butyl,  
35 (R)-1-(2-methyl)-butyl, cyclopentyl, 1-hexyl, (S)-2-hexyl, (R)-2-hexyl, (R)-3-hexyl, and



(S)-3-hexyl.

51. A compound according to claim 29, wherein  $R_4$ ,  $R_5$ , and  $R_6$ , are independently selected from hydrogen and methyl.


52. A compound according to claim 29, wherein any two of  $R_2$ ,  $R_4$ ,  $R_5$ , and  $R_6$  are hydrogen.

53. A compound according to claim 29 wherein  $d$  is 2.

54. A compound according to claim 53, wherein  $X$  is  $-S(O)_2-$ , and  $R_1$  is substituted or unsubstituted aralkyl.

55. A compound according to claim 54, wherein  $R_1$  is substituted or unsubstituted benzyl.

56. A compound of the formula:



(II)

wherein,

(a)  $X$  is selected from the group consisting of  $-S(O)_2-$ ,  $-N(R')-S(O)_2-$ ,  $-(C=O)-$ ,  $-OC(=O)-$ ,  $-NH-C(=O)-$ ,  $-P(O)(R'')$  and a direct link, wherein  $R'$  is hydrogen, alkyl of 1 to about 4 carbon atoms, aryl of about 6 to about 14 carbon atoms or aralkyl of about 6 to about 16 carbon atoms, and  $R''$  is  $-NR'$ ,  $-OR'$ ,  $-R'$ , or  $-SR'$ ;

(b)  $R_1$  is selected from the group consisting of:

- (1) alkyl of 1 to about 12 carbon atoms,
- (2) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 5 to about 8 carbon atoms, which optionally is substituted in the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',  
5
- (3) cycloalkyl of 3 to about 15 carbon atoms, which optionally is substituted in the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',  
10
- (4) heterocycloalkyl of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, wherein i is 0, 1, or 2, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',  
15
- (5) heterocyclo of 4 to about 10 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from the group consisting of oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',  
20
- (6) alkenyl of about 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8 carbon atoms, which optionally is substituted on the ring carbons with hydroxyl, amino, guanidino, amidino, C<sub>1</sub> to C<sub>3</sub> alkyl, C<sub>1</sub> to C<sub>3</sub> alkoxy, or CO<sub>2</sub>R',  
25
- (7) aryl of about 6 to about 14 carbon atoms which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,  
30
- (8) heteroaryl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>i</sub>, and which is optionally mono-, di- or tri-substituted on the  
35




ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>.


(9) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>,

(10) heteroaralkyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>1</sub>, which is optionally substituted on the alkyl chain with hydroxy or halogen and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,

(11) aralkenyl of about 8 to about 16 carbon atoms having 6 to about 14 ring carbons which is optionally mono-, di-, or tri-substituted on the aryl ring with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,

(12) heteroaralkenyl of 5 to 14 ring atoms with the ring atoms selected from carbon and heteroatoms, wherein the heteroatoms are selected from oxygen, nitrogen, and S(O)<sub>1</sub>, and which is optionally mono-, di- or tri-substituted on the ring carbons with Y<sub>1</sub>, Y<sub>2</sub>, and/or Y<sub>3</sub>, respectively,







(17) perfluoroalkyl of 1 to about 12 carbon atoms,

5 (18) perfluoroaryl of about 6 to about 14 carbon atoms,

(19) perfluoroaryl alkyl of about 7 to about 15 carbon atoms,

(20) hydrogen, and

10 (21) , wherein  is a 5 to 7 member heterocyclic ring having 3 to 6 ring carbon atoms, where V is  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{S}(=\text{O})-$ ,  $-\text{S}(\text{O})_2-$  or  $-\text{S}-$ , wherein  $\text{Y}_1$ ,  $\text{Y}_2$ , and  $\text{Y}_3$  are

(i) independently selected from the group  
 15 consisting of hydrogen, halogen, cyano, nitro, tetrazolyl, amino, guanidino, amidino, methylamino, methylguanidino,  $-\text{CF}_3$ ,  $-\text{CF}_2\text{CF}_3$ ,  $-\text{CF}_2\text{H}$ ,  $\text{CH}(\text{CF}_3)_2$ ,  $-\text{C}(\text{OH})(\text{CF}_3)_2$ ,  $-\text{OCF}_3$ ,  $-\text{OCF}_2\text{CF}_3$ ,  $-\text{OC}(\text{O})\text{NH}_2$ ,  $-\text{OC}(\text{O})\text{NHZ}_1$ ,  $-\text{OC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{NHC}(\text{O})\text{Z}_1$ ,  $-\text{NHC}(\text{O})\text{NH}_2$ ,  $-\text{NHC}(\text{O})\text{NZ}_1$ ,  $-\text{NHC}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OH}$ ,  $-\text{C}(\text{O})\text{NH}_2$ ,  
 20  $-\text{C}(\text{O})\text{NHZ}_1$ ,  $-\text{C}(\text{O})\text{NZ}_1\text{Z}_2$ ,  $-\text{C}(\text{O})\text{OZ}_1$ ,  $-\text{P}(\text{O})_3\text{H}$ ,  $-\text{P}(\text{O})_3\text{H}_2$ ,  $-\text{PH}(\text{O})\text{OH}$ ,  $-\text{P}(\text{O})_3(\text{Z}_1)_2$ ,  $-\text{S}(\text{O})_3\text{H}$ ,  $-\text{S}(\text{O})_m\text{Z}_1$ ,  $-\text{Z}_1$ ,  $-\text{OZ}_1$ ,  $-\text{OH}$ ,  $-\text{NH}_2$ ,  $-\text{NHZ}_1$ , and  $-\text{NZ}_1\text{Z}_2$ , wherein  $m$  is 0, 1 or 2, and  $\text{Z}_1$  and  $\text{Z}_2$  are independently selected from the group consisting of alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 14  
 25 carbon atoms, heteroaryl of about 5 to about 14 atoms having 1 to about 9 carbon atoms, aralkyl of about 7 to about 15 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms having about 3 to about 9 carbon atoms, or

(ii)  $\text{Y}_1$  and  $\text{Y}_2$  are selected together to be  
 30  $-\text{OC}[(\text{Z}_3)(\text{Z}_4)]_q\text{O}-$ , wherein  $q$  is an integer from 1 to 4 and  $\text{Z}_3$  and  $\text{Z}_4$  are independently selected from the group consisting of hydrogen, alkyl of 1 to about 12 carbon atoms, aryl of

about 6 to about 14 carbon atoms heter aryl f about 5 to about 14 atoms having 1 to about 9 carbon atoms, aralkyl of about 7 to about 15 carbon atoms, and heteroaralkyl of about 6 to about 11 atoms having about 3 to about 9 carbon atoms;

5 (c) R<sub>2</sub> is selected from the group consisting of hydrogen, alkyl of 1 to about 4 carbon atoms, and alkenyl of about 2 to about 4 carbon atoms;

(d) R<sub>4</sub> is selected from the group consisting of

(i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein n is 0, 1  
10 or 2, and R<sub>1</sub> is as defined above, with the proviso that R<sub>1</sub> is

not a camphor derivative or  heterocycle,

(ii) -CF<sub>3</sub>, -CF<sub>2</sub>H, -OCF<sub>3</sub>, -OCF<sub>2</sub>H,

(iii) halogen,

(iv) alkyl of 1 to 6 carbon atoms substituted with  
15 Z<sub>5</sub>, wherein Z<sub>5</sub> is selected from the group consisting of hydrogen, -OH, -OR<sub>8</sub>, -CH<sub>3</sub>, halogen, -C(O)OH, -C(O)OR<sub>8</sub> and -S(O)<sub>p</sub>R<sub>8</sub>, wherein R<sub>8</sub> is alkyl of 1 to about 6 carbon atoms, and p is 0, 1 or 2,

(v) alkyl of 1 to 3 carbon atoms substituted with  
20 cyclic alkyl of 3 to 5 carbon atoms,

(vi) alkenyl of about 3 to about 6 carbon atoms,

(vii) cycloalkyl of about 3 to about 10 carbon  
atoms,

(viii) heteroaryl of about 5 to about 10 ring atoms  
25 having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, and

(ix) heteroaralkyl of about 5 to about 10 ring  
atoms having 2 to about 15 carbon atoms which include 1 to  
30 about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen;

(e) R<sub>5</sub> is selected from the group consisting of

(i) -R<sub>1</sub>, -OR<sub>1</sub>, -NHR<sub>1</sub>, -S(O)<sub>n</sub>R<sub>1</sub>, wherein R<sub>1</sub> is as  
defined above, with the proviso that R<sub>1</sub> is not a camphor

derivative or  $\text{---}\overset{\text{N}}{\text{C}}\text{---}$  heter cycle,

(ii)  $\text{---CF}_3$ ,  $\text{---CF}_2\text{H}$ ,  $\text{---OCF}_3$ ,  $\text{---OCF}_2\text{H}$ , and

(iii) halogen;

(f)  $R_6$  is selected from the group consisting of

5 (i)  $\text{---R}_1$ ,  $\text{---OR}_1$ ,  $\text{---NHR}_1$ ,  $\text{---S(O)}_n\text{R}_1$ , wherein  $R_1$  is as defined above, with the proviso that  $R_1$  is not a camphor

derivative or  $\text{---}\overset{\text{N}}{\text{C}}\text{---}$  heterocycle,

(ii)  $\text{---CF}_3$ ,  $\text{---CF}_2\text{H}$ ,  $\text{---OCF}_3$ ,  $\text{---OCF}_2\text{H}$ ,

(iii) halogen,

10 (iv) alkyl of 1 to about 12 carbon atoms substituted with  $Z_6$ , wherein  $Z_6$  is selected from the group consisting of hydrogen, hydroxy, halogen,  $\text{---OR}_9$ ,  $\text{---NHR}_9$ ,  $\text{---C(O)OH}$ ,  $\text{---C(O)OR}_9$ , and  $\text{---S(O)}_p\text{R}_9$ , wherein  $R_9$  is alkyl of 1 to about 12 carbon atoms, aryl of about 6 to about 10 carbon  
15 atoms, aralkyl of about 7 to about 12 carbon atoms, heteroaryl of about 5 to about 10 ring atoms 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri- substituted in the ring  
20 carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ , and heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri-  
25 substituted in the ring carbons with  $Y_1$ ,  $Y_2$  and/or  $Y_3$ ,

(v) alkyl of 1 to about 3 carbon atoms substituted with cycloalkyl of about 3 to about 8 carbon atoms,

(vi) alkenyl of 2 to about 6 carbon atoms which is optionally substituted with cycloalkyl of about 3 to about 8  
30 carbon atoms, aryl of about 6 to about 10 carbon atoms, or heteroaryl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms,

(vii) aralkyl of about 7 to about 15 carbon atoms which is optionally substituted on the alkyl chain with

hydroxy or halogen, and optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(viii) heteroaralkyl of about 5 to about 10 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally substituted on the alkyl chain with hydroxy or halogen, and optionally mono-, di- or tri-substituted on the ring with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>,

(ix) aralkenyl of about 6 to 15 carbon atoms which is optionally mono-, di- or tri-substituted on the aryl with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>, and

(x) heteroaralkenyl ring of about 5 to about 10 ring atoms with 1 to 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, sulfur and nitrogen, which is optionally mono-, di- or tri-substituted with Y<sub>1</sub>, Y<sub>2</sub> and/or Y<sub>3</sub>;

(g) R<sub>7</sub> is selected from the group consisting of

(i) R<sub>1</sub>, wherein R<sub>1</sub> is as defined above, with the

proviso that R<sub>7</sub> is not an R<sub>1</sub> camphor derivative or heterocycle,


(ii) -CF<sub>3</sub> or CF<sub>2</sub>H,

(iii) alkyl of 1 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>, wherein each of R<sub>1</sub> and R<sub>11</sub> is independently selected and is alkyl of 1 to about 4 carbon atoms or hydrogen,

(iv) alkenyl of about 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, CONR<sub>10</sub>R<sub>11</sub>,

(v) cycloalkyl of 3 to about 10 carbon atoms, optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, -CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, -CONR<sub>10</sub>R<sub>11</sub>,

(vi) heteroaryl of about 5 to about 10 ring atoms, having about 1 to about 9 ring carbon atoms which is optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, -SO<sub>2</sub>R<sub>10</sub>, or



-CONR10R11, and

(vii) alkyl of 1 to about 3 carbon atoms substituted with heteroaryl of about 5 to about 16 atoms having 1 to about 9 carbon atoms, which is optionally substituted with -CH<sub>2</sub>OR<sub>10</sub>, CO<sub>2</sub>R<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, CONR<sub>10</sub>R<sub>11</sub>; and

(h) R<sub>12</sub> is selected from the group consisting of

(i) alkyl of 1 to about 12 carbon atoms which is optionally substituted with amino, amidino or guanidino,

(ii) aralkyl of 7 to about 15 carbon atoms which is optionally substituted on a ring carbon with amino, amidino, or guanidino,

(iii) aryl of about 6 to about 14 carbon atoms which is optionally substituted on a ring carbon with amino, amidino, or guanidino,

(iv) heteroaralkyl of 5 to about 14 ring atoms having 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur which is optionally substituted on a ring carbon with amino, amidino or guanidino; and

(j) TS is selected from the group consisting of

(i) -CHO,

(ii) -C(O)CF<sub>3</sub>,

(iii) -C(O)CF<sub>3</sub>CF<sub>3</sub>,

(iv) -C(O)-C(O)-NHR<sub>13</sub>,

(v) -C(O)-C(O)<sub>2</sub>R<sub>13</sub>,

(vi) -C(O)-C(O)-NR<sub>13</sub>R<sub>14</sub>,

(vii) -B(OR<sub>13</sub>)(OR<sub>14</sub>),

(viii) -C(O)CH<sub>2</sub>Cl, and

(ix) -C(O)-R<sub>13</sub>;


wherein R<sub>13</sub> and R<sub>14</sub> are independently selected from hydrogen, lower alkyl of 1 to about 6 carbon atoms, cycloalkyl of 3 to about 10 carbon atoms, aralkyl of 7 to about 15 carbon atoms, and heteroaralkyl of about 5 to about 10 ring atoms having about 2 to about 15 carbon atoms which include 1 to about 9 ring carbon atoms and the remainder of the ring atoms heteroatoms selected from oxygen, nitrogen and sulfur; or



when TS is  $-B(OR_{1,1})(OR_{1,2})$  then  $OR_{1,1}$  and  $OR_{1,2}$  may be taken together to be  $-O[C(Z_1)(Z_2)]qO-$ ; and  $R_{1,1}$  is selected from aryl of 6 to about 14 carbon atoms or heteroaryl of 5 to about 14 ring atoms having 1 to about 13 ring carbon atoms and the remainder of the ring atoms selected from oxygen, nitrogen and sulfur; and pharmaceutically acceptable salts thereof.

E

Figure 1



**Figure 2**




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7






Figure 8






Figure 9






Figure 10







Figure 11





Figure 12



# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/25167

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C279/12 C07C257/18 C07F5/04 C07D211/60

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 672 659 A (ELI LILLY AND COMPANY) 20 September 1995 see claim 1	1
A	EP 0 526 877 A (BRISTOL-MYERS SQUIBB COMPANY) 10 February 1993 see claim 1	1
A	FR 2 490 632 A (NIPPON KAYAKU KABUSHIKI KAISHA) 26 March 1982 see claim 1	1
A	WO 96 18644 A (CORVAS INTERNATIONAL, INC.) 20 June 1996 see claim 1	1
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

8 April 1999

Date of mailing of the international search report

16/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3018

Authorized officer

Kapteyn, H

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/25167

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 35313 A (CORVAS INTERNATIONAL, INC.) 28 December 1995 cited in the application see claim 1 -----	1
A	WO 95 35312 A (CORVAS INTERNATIONAL, INC.) 28 December 1995 see claim 1 -----	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/25167

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 672659 A	20-09-1995	US 5885967 A	23-03-1999
		AT 175404 T	15-01-1999
		CA 2143536 A	05-09-1995
		DE 69507062 D	18-02-1999
		JP 7258287 A	09-10-1995
EP 526877 A	10-02-1993	CA 2075154 A	07-02-1993
		JP 7242616 A	19-09-1995
		US 5380713 A	10-01-1995
FR 2490632 A	26-03-1982	JP 1585234 C	31-10-1990
		JP 2000342 B	08-01-1990
		JP 57054157 A	31-03-1982
		CA 1183130 A	26-02-1985
		DE 3137280 A	03-06-1982
		GB 2086380 A,B	12-05-1982
		US 4401594 A	30-08-1983
WO 9618644 A	20-06-1996	US 5656645 A	12-08-1997
		US 5658930 A	19-08-1997
		AU 693636 B	02-07-1998
		AU 4424896 A	03-07-1996
		BR 9509994 A	30-12-1997
		CA 2206400 A	20-06-1996
		EP 0804464 A	05-11-1997
		HU 77888 A	28-09-1998
		JP 10510539 T	13-10-1997
WO 9535313 A	28-12-1995	US 5714499 A	03-02-1998
		AT 176241 T	15-02-1999
		AU 2905495 A	15-01-1996
		CA 2192211 A	28-12-1995
		DE 69507614 D	11-03-1999
		EP 0765339 A	02-04-1997
		JP 10503177 T	24-03-1998
		AU 700808 B	14-01-1999
		AU 2863095 A	15-01-1996
		BR 9508048 A	18-11-1997
		CA 2192210 A	28-12-1995
		CN 1151166 A	04-06-1997
		EP 0802923 A	29-10-1997
		JP 10503176 T	24-03-1998
		NO 965353 A	17-02-1997
		NZ 288940 A	28-01-1999
		WO 9535311 A	28-12-1995
		US 5703208 A	30-12-1997
WO 9535312 A	28-12-1995	US 5637599 A	10-06-1997
		AU 2905095 A	15-01-1996
		CA 2192686 A	28-12-1995
		EP 0765338 A	02-04-1997
		JP 10506094 T	16-06-1998
		US 5883107 A	16-03-1999